

=> d his

(FILE 'HOME' ENTERED AT 12:23:26 ON 03 JUN 2002)

FILE 'REGISTRY' ENTERED AT 12:24:19 ON 03 JUN 2002

L1 1 S 9004-54-0 ← CMD  
 L2 921 S 9004-54-0/CRN ← all mixtures (registered) comprising  
 L3 1 S 79-14-1 ← 170-CH<sub>2</sub>-O-17  
 L4 1936 S 79-14-1/CRN ← mixtures containing  
 L5 70 S L2 AND L4  
 L6 509625 S OC5/ES ← any pyran  
 L7 4 S L6 AND L5  
 L8 216531 S PETH/PCT ← any polymer w/ a poly ether  
 L9 2775 S L6 AND L8  
 L10 1216 S L9 AND "2-ETHANEDIYL" cpds w/ CH<sub>2</sub>-CH<sub>2</sub>-O- frag  
 L11 0 S L2 AND L10  
 L12 27 S L2 AND L8  
 L13 0 S L9 AND L2  
 L14 2 S L4 AND L12  
 L15 1 S 195214-71-2

FILE 'HCAPLUS' ENTERED AT 12:34:33 ON 03 JUN 2002

L16 435 S L5  
 L17 470851 S GLUCOSE OR GALACTOSE OR HEXOSE OR ?SACCHARIDE  
 L18 14 S L16(L)L17  
 L19 304673 S POLYETHYLEN? OR PEG OR ?ETHANEDIYL?  
 L20 13059 S LINKER  
 L21 262 S L19(L)L20  
 L22 0 S L21 AND L18  
 L23 1 S L18 AND L19  
 L24 1 S L18 AND L20  
 L25 2 S L23-24  
 L26 98 S L16 AND L17  
 L27 10 S L26 AND L19  
 L28 3 S L26 AND L20  
 L29 13 S L26 AND L27-28  
 L30 10 S L29 NOT L28  
 L31 7249 S L17(L)?CONJUGAT?  
 L32 29 S L31 AND L16  
 L33 3 S L32 AND (L19 OR L20)  
 L34 1042 S ?CARBOXYMETHYL(2W)DEXTRAN? OR CMD  
 L35 1099 S ?CARBOXYMETHYL?(2W)?DEXTRAN? OR CMD  
 L36 57 S L35 NOT L34  
 L37 71 S L34(L)CONJUGAT?  
 L38 18 S L31 AND L37  
 L39 3 S L28 AND L19-20  
 L40 18 S L38 NOT L39  
 L41 22 S L39-40 OR L33 OR L28 OR L25  
 SELECT RN L41 1-22

FILE 'REGISTRY' ENTERED AT 12:55:07 ON 03 JUN 2002

L42 200 S E1-200  
 L43 46 S E201-246  
 L44 246 S L42-43  
 L45 28 S L44 AND L1-15  
 L46 218 S L44 NOT L45

FILE 'HCAPLUS' ENTERED AT 12:58:28 ON 03 JUN 2002

L47 22 S L41 AND L44  
 22 citations

=> d ibib abs hitstr 1-22

L47 ANSWER 1 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2002:148739 HCAPLUS

DOCUMENT NUMBER: 136:205403

TITLE: DDS compounds of drugs having hydroxy groups

INVENTOR(S): Ousu, Satoru; Oki, Hitoshi; Naito, Hiroyuki; Hirotsu, Kenji

PATENT ASSIGNEE(S): Daiichi Seiyaku Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 24 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2002060351	A2	20020226	JP 2001-80188	20010321

PRIORITY APPLN. INFO.: JP 2000-79655 A 20000322

OTHER SOURCE(S): MARPAT 136:205403

AB The DDS (drug delivery system) compds. are represented by the formula  
 AWN(R1)C(R2)(R3)OQ or PZN(R1)C(R2)(R3)OQ [A = polymeric carrier for drugs;  
 W = spacer contg. amino acid or oligopeptide residue linked to N at the  
 C-terminal; P = protective group for H or NH<sub>2</sub>; Z = amino acid residue or  
 oligopeptide residue linked to N at the C-terminal; R1-R3 = H,  
 (substituted) alkyl, (substituted) aryl, carboxyl, alkoxy-carbonyl; 2 of  
 R1-R3 may form 4- to 8-membered ring; OQ = residue of OH-contg. drugs].  
 Tert-Bu 13-[[1-[2-amino-6-[4-[(E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-  
 1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy]-7-  
 benzyl-2,5,8,11-tetraoxo-3,6,9,12-tetraazatri-1-decylcarbamate (prepn.  
 given) showed 89% release of 1-[2-amino-6-[4-[(E)-3-[4-(3,5-  
 difluorophenyl)-1-piperazinyl]-1-propenyl]-1H-pyrazol-1-yl]-4-pyrimidinyl]-  
 3-azetidinol (I) in murine fibrosarcoma Meth-A cell homogenate at  
 40.degree. and pH 4.5 and <1% release of I in a buffer at pH 4.5. I.v.  
 administration of a carboxymethyl dextran polyol deriv. of I (linked  
 through an oligopeptide and aminomethylene linker) at 10 mg/kg  
 as I showed significant antitumor effect and did not cause diarrhea in  
 mice.

IT 401470-57-3


RL: PAC (Pharmacological activity); RCT (Reactant); THU (Therapeutic use);  
 BIOL (Biological study); RACT (Reactant or reagent); USES (Uses)

(prepn. of amino acid or peptide derivs. of hydroxy-contg. drugs for  
 DDS)

RN 401470-57-3 HCAPLUS

CN 3-Azetidinol, 1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-  
 piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]- (9CI)  
 (CA INDEX NAME)


Double bond geometry as shown.



IT 39422-83-8DP, Carboxymethyldextran sodium salt, polyols,  
 conjugates with peptide spacers and antitumor drugs 401470-32-4P  
 401470-34-6DP, conjugates with carboxymethyl dextran polyols  
 401470-36-8DP, conjugates with carboxymethyl dextran polyols  
 401470-38-0DP, conjugates with carboxymethyl dextran polyols  
 401470-40-4DP, conjugates with carboxymethyl dextran polyols  
 401470-44-8DP, conjugates with carboxymethyl dextran polyols  
 401470-48-2DP, conjugates with carboxymethyl dextran polyols  
 401470-52-8DP, conjugates with carboxymethyl dextran polyols  
 401470-56-2DP, conjugates with carboxymethyl dextran polyols  
 RL: PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use);  
 BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (prepn. of amino acid or peptide derivs. of hydroxy-contg. drugs for  
 DDS)  
 RN 39422-83-8 HCAPLUS  
 CN Dextran, carboxymethyl ether, sodium salt (9CI) (CA INDEX NAME)  
 CM 1  
 CRN 9004-54-0  
 CMF Unspecified  
 CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE


CM 2  
 CRN 79-14-1  
 CMF C2 H4 O3




RN 401470-32-4 HCAPLUS  
 CN Glycinamide, N-[(1,1-dimethylethoxy)carbonyl]glycylglycyl-L-phenylalanyl-N-  
 [[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-  
 propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-  
 azetidinyloxy)methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
 Double bond geometry as shown.

PAGE 1-A







RN 401470-34-6 HCAPLUS

CN Glycinamide, L-phenylalanyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy)methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.




RN 401470-36-8 HCAPLUS


CN Glycinamide, glycyglycyl-L-phenylalanyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy)methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A



PAGE 1-B




RN 401470-38-0 HCAPLUS


CN Glycinamide, glycyglycyl-L-isoleucyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy]methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A



PAGE 1-B




RUSSEL 09/807,980

RN 401470-40-4 HCAPLUS


CN Glycinamide, glycyglycyl-L-valyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy)methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A



PAGE 1-B





RN 401470-44-8 HCAPLUS

CN Glycinamide, glycyglycyl-L-methionyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy)methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A







RN 401470-48-2 HCAPLUS

CN Glycinamide, glycyglycyl-L-phenylalanyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy]methyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.




RN 401470-52-8 HCAPLUS


CN L-Isoleucinamide, glycyglycyl-L-prolyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy]methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A



PAGE 1-B




RN 401470-56-2 HCAPLUS


CN L-Leucinamide, glycyglycyl-L-prolyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy]methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A








IT 16257-05-9 18621-17-5, 1-Benzhydryl-3-azetidinol  
 31972-52-8 32991-17-6 35661-40-6  
 35665-38-4 39621-73-3 71989-28-1  
 160036-44-2 256930-32-2 333366-34-0  
 401470-59-5

RL: RCT (Reactant); RACT (Reactant or reagent)  
 (prepn. of amino acid or peptide derivs. of hydroxy-contg. drugs for  
 DDS)

RN 16257-05-9 HCAPLUS


CN Glycine, N-[(1,1-dimethylethoxy)carbonyl]-L-isoleucyl- (9CI) (CA INDEX  
 NAME)

Absolute stereochemistry.




RN 18621-17-5 HCAPLUS

CN 3-Azetidinol, 1-(diphenylmethyl)- (8CI, 9CI) (CA INDEX NAME)



RN 31972-52-8 HCAPLUS


CN Glycine, N-[(1,1-dimethylethoxy)carbonyl]glycyl- (9CI) (CA INDEX NAME)



RN 32991-17-6 HCAPLUS


CN Glycine, N-[(1,1-dimethylethoxy)carbonyl]-L-leucyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 35661-40-6 HCAPLUS  
CN L-Phenylalanine, N-[(9H-fluoren-9-ylmethoxy)carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




RN 35665-38-4 HCAPLUS  
CN Glycine, N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycyl- (9CI) (CA INDEX NAME)




RN 39621-73-3 HCAPLUS  
CN L-Phenylalanine, N-[(1,1-dimethylethoxy)carbonyl]glycylglycyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 71989-28-1 HCAPLUS  
CN L-Methionine, N-[(9H-fluoren-9-ylmethoxy)carbonyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 160036-44-2 HCAPLUS

CN L-Phenylalanine, N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycylglycyl- (9CI)  
(CA INDEX NAME)


Absolute stereochemistry.



RN 256930-32-2 HCAPLUS

CN 2-Pyrimidinamine, 4-chloro-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]- (9CI) (CA INDEX NAME)


Double bond geometry as shown.



RN 333366-34-0 HCAPLUS

CN L-Proline, N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycylglycyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 401470-59-5 HCAPLUS

CN L-Isoleucine, N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycylglycyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 60667-52-9P 401470-30-2P 401470-31-3P  
 401470-33-5P 401470-34-6P 401470-35-7P  
 401470-36-8P 401470-37-9P 401470-38-0P  
 401470-39-1P 401470-40-4P 401470-41-5P  
 401470-42-6P 401470-43-7P 401470-44-8P  
 401470-45-9P 401470-46-0P 401470-47-1P  
 401470-48-2P 401470-49-3P 401470-50-6P  
 401470-51-7P 401470-52-8P 401470-53-9P  
 401470-54-0P 401470-55-1P 401470-56-2P  
 401470-58-4P 401470-60-8P 401470-61-9P  
 401470-62-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(prepn. of amino acid or peptide derivs. of hydroxy-contg. drugs for DDS)

RN 60667-52-9 HCAPLUS

CN Glycine, N-[(1,1-dimethylethoxy)carbonyl]-N-methylglycyl- (9CI) (CA INDEX NAME)




RN 401470-30-2 HCAPLUS


CN Carbamic acid, [2-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy)methyl]amino]-2-oxoethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Double bond geometry as shown.

PAGE 1-A




PAGE 1-B



RN 401470-31-3 HCAPLUS

CN Acetamide, N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy]methyl]- (9CI) (CA INDEX NAME)

Double bond geometry as shown.





RN 401470-33-5 HCAPLUS -

CN Glycinamide, N-[(9H-fluoren-9-ylmethoxy)carbonyl]-L-phenylalanyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy]methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Double bond geometry as shown.




RN 401470-34-6 HCAPLUS

CN Glycinamide, L-phenylalanyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy]methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.







RN 401470-35-7 HCAPLUS

CN Glycinamide, N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycylglycyl-L-phenylalanyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy)methyl]- (9CI) (CA INDEX NAME)



Absolute stereochemistry.  
Double bond geometry as shown.



RN 401470-36-8 HCAPLUS

CN Glycinamide, glycylglycyl-L-phenylalanyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy)methyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.  
Double bond geometry as shown.




RN 401470-37-9 HCAPLUS

CN Glycinamide, N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycylglycyl-L-isoleucyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy]methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.









RN 401470-38-0 HCAPLUS

CN Glycinamide, glycyglycyl-L-isoleucyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy]methyl]- (9CI) (CA INDEX NAME)



Absolute stereochemistry.  
Double bond geometry as shown.



RN 401470-39-1 HCAPLUS

CN Glycinamide, N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycyglycyl-L-valyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy]methyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.  
Double bond geometry as shown.




RN 401470-40-4 HCAPLUS

CN Glycinamide, glycylglycyl-L-valyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy]methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.







RN 401470-41-5 HCAPLUS

CN Glycinamide, N-[(9H-fluoren-9-ylmethoxy)carbonyl]-L-methionyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy]methyl]- (9CI)  
(CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.




RN 401470-42-6 HCAPLUS


CN Glycinamide, L-methionyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy]methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A



PAGE 1-B





RN 401470-43-7 HCAPLUS

CN Glycinamide, N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycylglycyl-L-methionyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy]methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A







RN 401470-44-8 HCAPLUS

CN Glycinamide, glycylglycyl-L-methionyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy)methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.




RN 401470-45-9 HCAPLUS


CN Carbamic acid, [2-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy)methyl]amino]-2-oxoethyl]methyl-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Double bond geometry as shown.

PAGE 1-A



PAGE 1-B




RN 401470-46-0 HCAPLUS


CN Acetamide, N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy]methyl]-2-(methylamino)- (9CI) (CA INDEX NAME)

Double bond geometry as shown.

PAGE 1-A



PAGE 1-B




RN 401470-47-1 HCAPLUS

CN Glycinamide, N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycylglycyl-L-phenylalanyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy]methyl]-2-(methylamino)- (9CI) (CA INDEX NAME)


azetidinyloxy)methyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A



PAGE 1-B





RN 401470-48-2 HCAPLUS

CN Glycinamide, glycyglycyl-L-phenylalanyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy)methyl]-N2-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A







RN 401470-49-3 HCAPLUS

CN Carbamic acid, [(1S,2S)-1-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy]methyl]amino]carbonyl]-2-methylbutyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.





RN 401470-50-6 HCAPLUS

CN Pentanamide, 2-amino-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy]methyl]-3-methyl-, (2S,3S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.







RN 401470-51-7 HCAPLUS

CN L-Isoleucinamide, N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycylglycyl-L-prolyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy]methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.





NH<sub>2</sub>



RN 401470-52-8 HCAPLUS

CN L-Isoleucinamide, glycylglycyl-L-prolyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy)methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.



RN 401470-53-9 HCAPLUS


CN Carbamic acid, [(1S)-1-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyloxy)methyl]amino]carbonyl]-3-methylbutyl]-,

RUSSEL 09/807,980


1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A



PAGE 1-B





RN 401470-54-0 HCAPLUS

CN Pentanamide, 2-amino-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-azetidinyl]oxy)methyl]-4-methyl-, (2S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A






RN 401470-55-1 HCAPLUS


CN L-Leucinamide, N-[(9H-fluoren-9-ylmethoxy)carbonyl]glycylglycyl-L-prolyl-N-  
[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-difluorophenyl)-1-piperazinyl]-1-  
propenyl]-5-methyl-1H-pyrazol-1-yl]-4-pyrimidinyl]-3-  
azetidinyloxy)methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A



PAGE 1-B




RN 401470-56-2 HCAPLUS


CN L-Leucinamide, glycylglycyl-L-prolyl-N-[[[1-[2-amino-6-[4-[(1E)-3-[4-(3,5-  
difluorophenyl)-1-piperazinyl]-1-propenyl]-5-methyl-1H-pyrazol-1-yl]-4-  
pyrimidinyl]-3-azetidinyloxy)methyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.  
Double bond geometry as shown.

PAGE 1-A




PAGE 1-B




RN 401470-58-4 HCAPLUS

CN Carbamic acid, [2-[[[1-(diphenylmethyl)-3-azetidinyl]oxy]methyl]amino]-2-oxoethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)



RN 401470-60-8 HCAPLUS


CN Carbamic acid, [2-[[[1-(diphenylmethyl)-3-azetidinyl]oxy]methyl]amino]-2-oxoethyl]methyl-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)



RN 401470-61-9 HCAPLUS

CN Carbamic acid, [(1S,2S)-1-[[[1-(diphenylmethyl)-3-azetidinyl]oxy]methyl]amino]carbonyl]-2-methylbutyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 401470-62-0 HCAPLUS

CN Carbamic acid, [(1S)-1-[[[1-(diphenylmethyl)-3-azetidinyloxy]methyl]amino]carbonyl]-3-methylbutyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L47 ANSWER 2 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2001:669315 HCAPLUS

DOCUMENT NUMBER: 136:18928

TITLE: New antibody purification procedure using a thermally responsive poly(N-isopropylacrylamide)-dextran derivative conjugate

AUTHOR(S): Anastase-Ravion, S.; Ding, Z.; Pelle, A.; Hoffman, A. S.; Letourneur, D.

CORPORATE SOURCE: INVIMAT, Universite Paris 13, Villetaneuse, 93430, Fr.

SOURCE: Journal of Chromatography, B: Biomedical Sciences and Applications (2001), 761(2), 247-254  
CODEN: JCBBEP; ISSN: 0378-4347

PUBLISHER: Elsevier Science B.V.


DOCUMENT TYPE: Journal

LANGUAGE: English


AB Through their specificity and affinity, antibodies are useful tools in research and medicine. In this study, we investigated a new type of chromatog. method using a thermosensitive polymer for the purifn. of antibodies against a dextran deriv. (DD), as a model. The thermally reversible sol.-insol. poly(N-isopropylacrylamide)-dextran deriv. **conjugate**, named poly(NIPAAm)-DD, has been synthesized by **conjugating** amino-terminated poly(N-isopropylacrylamide) to a DD via ethyl-3-(3-dimethylaminopropyl)-carbodiimide. On one hand, this report describes the two steps of poly(NIPAAm)-DD **conjugation** and characterization. On the other hand, the poly(NIPAAm)-DD **conjugate** was used as a tool to purify polyclonal antibodies in serum samples from rabbits s.c. immunized with the derivatized dextran. Antibodies were purified and quantified by immunoenzymic assays. Our results indicate that antibodies recognized both DD and poly(NIPAAm)-DD. In contrast, they did not bind to native poly(NIPAAm) or poly(NIPAAm) **conjugated** with another anionic dextran. We conclude that the

**conjugation** of a **polysaccharide** to poly(NIPAAm) leads to an original and efficient chromatog. method to purify antibodies. Moreover, this novel method of purifn. is rapid, sensitive, inexpensive and could be used to purify various types of antibodies.


IT **25189-55-3**, Poly(N-isopropylacrylamide)  
 RL: PRP (Properties); RCT (Reactant); RACT (Reactant or reagent)  
 (new antibody purifn. procedure using a thermally responsive  
 poly(N-isopropylacrylamide)-dextran deriv. conjugate)  
 RN 25189-55-3 HCAPLUS  
 CN 2-Propenamide, N-(1-methylethyl)-, homopolymer (9CI) (CA INDEX NAME)  
 CM 1  
 CRN 2210-25-5  
 CMF C6 H11 N O



IT **156-57-0**, 2-Aminoethanethiol hydrochloride **66205-07-0**,  
 2,2'-Azobisbutyronitrile  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (new antibody purifn. procedure using a thermally responsive  
 poly(N-isopropylacrylamide)-dextran deriv. conjugate)  
 RN 156-57-0 HCAPLUS  
 CN Ethanethiol, 2-amino-, hydrochloride (8CI, 9CI) (CA INDEX NAME)



RN 66205-07-0 HCAPLUS  
 CN Butanenitrile, 2,2'-azobis- (9CI) (CA INDEX NAME)




IT **9044-05-7DP**, Carboxymethyl dextran,  
 poly(N-isopropylacrylamide) **conjugates** 25189-55-3DP,  
 Poly(N-isopropylacrylamide), **carboxymethyl dextran**  
**conjugates**  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)  
 (new antibody purifn. procedure using a thermally responsive  
 poly(N-isopropylacrylamide)-dextran deriv. **conjugate**)  
 RN 9044-05-7 HCAPLUS  
 CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)  
 CM 1

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2


CRN 79-14-1  
CMF C2 H4 O3



RN 25189-55-3 HCAPLUS  
CN 2-Propenamide, N-(1-methylethyl)-, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 2210-25-5  
CMF C6 H11 N O



REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 3 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2001:322648 HCAPLUS

DOCUMENT NUMBER: 135:185307

TITLE: Characteristics of tissue distribution of various polysaccharides as drug carriers: influences of molecular weight and anionic charge on tumor targeting  
AUTHOR(S): Sugahara, Shuichi; Okuno, Satoshi; Yano, Toshiro; Hamana, Hiroshi; Inoue, Kazuhiro

CORPORATE SOURCE: Drug Delivery System Institute, Ltd., Chiba, 278-0022, Japan

SOURCE: Biological & Pharmaceutical Bulletin (2001), 24(5), 535-543

CODEN: BPBLEO; ISSN: 0918-6158

PUBLISHER: Pharmaceutical Society of Japan

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Using the Walker 256 model for carcinosarcoma-bearing rats, we i.v. administered 5 **polysaccharide** carriers with various mol. wts. (MWs) and elec. charges and tested for their plasma and tissue distribution. Two carriers, carboxymethylated-D-manno-D-glucan (CMMG) and CMDextran (CMDex), showed higher plasma AUC than the other carriers tested, namely, CMchitin (CMCh), N-desulfated N-acetylated heparin (DSH), and hyaluronic acid (HA). This was consistently found to be true over the range of MWs tested. For CMDex, the max. value of plasma AUC was obtained when the MW exceeded 150 kDa. As for the anionic charge, CMDex (110-180 kDa) with a degree of substitution (DS) of the CM groups ranging from 0.2 to 0.6, showed max. plasma AUC values. Twenty-four hours after administration, the concn. of CMDex (180-250 kDa; DS: 0.6-1.2) in tumors



was more than 3% of dose/g-approx. 10-fold higher than those obsd. with CMCh, DSH and HA. Doxorubicin (DXR) was bound to these carriers via a peptide spacer, GlyGlyPheGly (GGFG), to give carrier-GGFG-DXR **conjugates** (DXR content: 4.2-7.0 (wt./wt.)), and the antitumor effects of these **conjugates** were tested with Walker 256 carcinosarcoma-bearing rats by monitoring the tumor wts. after a single i.v. injection. Compared with free DXR, CMDex-GGFG-DXR and CMMG-GGFG-DXR **conjugates** significantly suppressed tumor growth, while the CMCh-GGFG-DXR, DSH-GGFG-DXR, and HA-GGFG-DXR **conjugates** in a similar comparison showed weak tumor growth inhibition. These findings suggest that the antitumor effect of the carrier-DXR **conjugates** was related to the extent with which the carriers accumulated in the tumors.

IT 9067-32-7DP, Hyaluronic acid sodium salt, **conjugates** with doxorubicin and peptide 23214-92-8DP, Doxorubicin, **conjugates** with peptide and polysaccharides 39422-83-8DP, Carboxymethyl dextran sodium salt, **conjugates** with doxorubicin and peptide 65667-26-7DP, **conjugates** with doxorubicin and peptide 105156-94-3DP, Carboxymethyl chitin sodium salt, **conjugates** with doxorubicin and peptide 200427-88-9DP, **conjugates** with doxorubicin and polysaccharides 355129-33-8DP, **conjugates** with doxorubicin and peptide  
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)  
 (effects of mol. wt. and anionic charge of **polysaccharide** drug carriers on tumor targeting)

RN 9067-32-7 HCAPLUS


CN Hyaluronic acid, sodium salt (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 23214-92-8 HCAPLUS

CN 5,12-Naphthacenedione, 10-[(3-amino-2,3,6-trideoxy-.alpha.-L-lyxo-hexopyranosyl)oxy]-7,8,9,10-tetrahydro-6,8,11-trihydroxy-8-(hydroxyacetyl)-1-methoxy-, (8S,10S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 39422-83-8 HCAPLUS

CN Dextran, carboxymethyl ether, sodium salt (9CI) (CA INDEX NAME)

CM 1

CRN 9004-54-0


CMF Unspecified

CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
CMF C2 H4 O3



RN 65667-26-7 HCAPLUS  
CN Heparamine, N-acetyl, sodium salt (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 105156-94-3 HCAPLUS  
CN Chitin, carboxymethyl ether, sodium salt (9CI) (CA INDEX NAME)


CM 1

CRN 1398-61-4  
CMF Unspecified  
CCI MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE


CM 2

CRN 79-14-1  
CMF C2 H4 O3



RN 200427-88-9 HCAPLUS  
CN Glycine, glycylglycyl-L-phenylalanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 355129-33-8 HCAPLUS  
CN D-Gluc-D-mannan, carboxymethyl ether, sodium salt (9CI) (CA INDEX NAME)

CM 1


CRN 11078-31-2  
CMF Unspecified

CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
CMF C2 H4 O3



IT 9067-32-7P, Hyaluronic acid sodium salt 39422-83-8P,  
Carboxymethyl dextran sodium salt 65667-26-7P  
105156-94-3P, Carboxymethyl chitin sodium salt  
355129-33-8P

RL: PRP (Properties); RCT (Reactant); SPN (Synthetic preparation); THU  
(Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT  
(Reactant or reagent); USES (Uses)  
(effects of mol. wt. and anionic charge of polysaccharide drug carriers  
on tumor targeting)

RN 9067-32-7 HCAPLUS

CN Hyaluronic acid, sodium salt (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 39422-83-8 HCAPLUS

CN Dextran, carboxymethyl ether, sodium salt (9CI) (CA INDEX NAME)


CM 1

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
CMF C2 H4 O3



RN 65667-26-7 HCAPLUS

CN Heparamine, N-acetyl, sodium salt (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 105156-94-3 HCAPLUS

CN Chitin, carboxymethyl ether, sodium salt (9CI) (CA INDEX NAME)


CM 1

CRN 1398-61-4  
CMF Unspecified  
CCI MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
CMF C2 H4 O3



RN 355129-33-8 HCAPLUS  
CN D-Gluco-D-mannan, carboxymethyl ether, sodium salt (9CI) (CA INDEX NAME)


CM 1

CRN 11078-31-2  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
CMF C2 H4 O3



REFERENCE COUNT: 43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 4 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 2000:228007 HCAPLUS

DOCUMENT NUMBER: 133:109911

TITLE: Ionic **Polysaccharide** Hydrogels via the  
Passerini and Ugi Multicomponent Condensations:  
Synthesis, Behavior and Solid-State NMR  
Characterization

AUTHOR(S): De Nooy, Arjan E. J.; Capitani, Donatella; Masci,  
Giancarlo; Crescenzi, Vittorio

CORPORATE SOURCE: Department of Chemistry, University 'La Sapienza',  
Rome, 00185, Italy

SOURCE: Biomacromolecules (2000), 1(2), 259-267  
CODEN: BOMAF6; ISSN: 1525-7797

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Original data are provided demonstrating that the title condensations are simple and versatile methods for the synthesis of hydrogels based on a variety of carboxylated polysaccharides. In this work, the biopolymers considered are sodium hyaluronate and sodium alginate. Nonnatural carboxylated polysaccharides were com. (carboxymethyl)cellulose or were obtained by carboxymethylation or selective oxidn. of primary alc. groups of scleroglucan and dextran. Hydrogels prepd. via the Passerini reaction

were transparent, alkali labile materials whereas the transparency of the Ugi gels depended on the **polysaccharide**, the cross-linker, and the degree of crosslinking. The Ugi gels were stable for several months at a pH ranging from 1.3 to 11 and up to temps. over 90 .degree.C. The structure of the networks was studied by means of <sup>13</sup>C CP-MAS and <sup>15</sup>N CP-MAS NMR spectroscopy. A quant. NMR anal. and elemental anal. of the dry gels allowed us to est. the efficiency of the reactions, i.e., the actual degree of crosslinking, which appeared to be about 80% of theor. The influence of added salt and pH on the swelling of several Ugi gels with different degrees of crosslinking was studied in a qual. manner.


IT 462-94-2, 1,5-Pentanediamine 931-53-3, Cyclohexyl isocyanide 2769-64-4, Butyl isocyanide 4117-33-3, L-Lysine ethyl ester 9004-32-4 9005-38-3, Sodium alginate 9044-05-7, Carboxymethyl dextran 9067-32-7, Sodium hyaluronate 39464-87-4D, Scleroglucan, oxidized 282730-55-6

RL: PEP (Physical, engineering or chemical process); PRP (Properties); RCT (Reactant); THU (Therapeutic use); BIOL (Biological study); PROC (Process); RACT (Reactant or reagent); USES (Uses)

(prepn., behavior and solid-state NMR characterization of ionic **polysaccharide** hydrogels via the Passerini and Ugi multicomponent condensations)


RN 462-94-2 HCAPLUS

CN 1,5-Pentanediamine (8CI, 9CI) (CA INDEX NAME)




RN 931-53-3 HCAPLUS

CN Cyclohexane, isocyano- (9CI) (CA INDEX NAME)



RN 2769-64-4 HCAPLUS


CN Butane, 1-isocyano- (9CI) (CA INDEX NAME)



RN 4117-33-3 HCAPLUS

CN L-Lysine, ethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 9004-32-4 HCAPLUS

CN Cellulose, carboxymethyl ether, sodium salt (8CI, 9CI) (CA INDEX NAME)

CM 1  
CRN 9004-34-6  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2  
CRN 79-14-1  
CMF C2 H4 O3



RN 9005-38-3 HCAPLUS  
CN Alginic acid, sodium salt (8CI, 9CI) (CA INDEX NAME)


\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9044-05-7 HCAPLUS  
CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1  
CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2  
CRN 79-14-1  
CMF C2 H4 O3



RN 9067-32-7 HCAPLUS  
CN Hyaluronic acid, sodium salt (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 39464-87-4 HCAPLUS  
CN Scleroglucan (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 282730-55-6 HCAPLUS  
CN Scleroglucan, carboxymethyl ether (9CI) (CA INDEX NAME)


CM 1  
CRN 39464-87-4  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1

CMF C2 H4 O3



REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 5 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1999:684285 HCAPLUS

DOCUMENT NUMBER: 131:269259

TITLE: Affinity-type biosensor with gold surface, **linker** layer and hydrogel, and method for the fabrication

INVENTOR(S): Wischerhoff, Erik; Nicolaus, Thomas

PATENT ASSIGNEE(S): BioTul Bio Instruments G.m.b.H., Germany

SOURCE: Ger. Offen., 12 pp.

CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 19817180	A1	19991021	DE 1998-19817180	19980417
DE 19817180	C2	20000427		

AB The invention concerns an affinity biosensor and its fabrication, that is composed of a gold surface, **linker** mols. and a bound hydrogel layer; the coverage of the surface is completed via hydrogen bonds and interactions of arom. rings upon sol-gel transformation. The **linker** mols. are of the general formula ARB; A = gold binding group, e.g. thio, disulfide, selenide, ; R = hydrocarbon chain, contg. at least two isolated phenol groups or heteroatoms; B = hydrogen binding group, e.g. hydroxy, epoxy, amino. Hydrogels are **polysaccharide** derivs., e.g. carboxymethyldextran. Alternately, the hydrogel is bound via a metal oxide layer to the **linker**.


IT 60-24-2, Mercaptoethanol 2602-34-8, Silane, [3-(2,3-epoxypropoxy)propyl]triethoxy- 5593-70-4 6066-82-6, N-Hydroxysuccinimide 7440-57-5, Gold, uses 9044-05-7, Carboxymethyldextran 17173-68-1, Ethanamine, 2,2'-dithiobis-, hydrochloride 25952-53-8, 1,3-Propanediamine, N'-(ethylcarbonimidoyl)-N,N-dimethyl-, monohydrochloride RL: DEV (Device component use); PEP (Physical, engineering or chemical process); PROC (Process); USES (Uses) (affinity-type biosensor with gold surface, **linker** layer and hydrogel, and method for fabrication)

RN 60-24-2 HCAPLUS



CN Ethanol, 2-mercapto- (8CI, 9CI) (CA INDEX NAME)




RN 2602-34-8 HCAPLUS  
CN Silane, triethoxy[3-(oxiranylmethoxy)propyl]- (9CI) (CA INDEX NAME)



RN 5593-70-4 HCAPLUS  
CN 1-Butanol, titanium(4+) salt (9CI) (CA INDEX NAME)



RN 6066-82-6 HCAPLUS  
CN 2,5-Pyrrolidinedione, 1-hydroxy- (9CI) (CA INDEX NAME)



RN 7440-57-5 HCAPLUS  
CN Gold (8CI, 9CI) (CA INDEX NAME)

Au

RN 9044-05-7 HCAPLUS  
CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)


CM 1

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE


CM 2

CRN 79-14-1  
CMF C2 H4 O3





RN 17173-68-1 HCAPLUS  
CN Ethanamine, 2,2'-dithiobis-, hydrochloride (9CI) (CA INDEX NAME)



●x HCl

RN 25952-53-8 HCAPLUS  
CN 1,3-Propanediamine, N'-(ethylcarbonimidoyl)-N,N-dimethyl-,  
monohydrochloride (9CI) (CA INDEX NAME)



● HCl

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 6 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1999:9887 HCAPLUS

DOCUMENT NUMBER: 130:71612

TITLE: Bioresorbable antiadhesion of carboxypolysaccharide  
polyether intermacromolecular complexes and methods  
for their use in reducing surgical adhesions

INVENTOR(S): Schwartz, Herbert E.; Blackmore, John M.

PATENT ASSIGNEE(S): Fziomed, Inc., USA

SOURCE: PCT Int. Appl., 95 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9858011	A1	19981223	WO 1998-US10814	19980528
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
US 5906997	A	19990525	US 1997-877649	19970617
US 6017301	A	20000125	US 1998-23267	19980213
US 6034140	A	20000307	US 1998-23097	19980213
AU 9876985	A1	19990104	AU 1998-76985	19980528
EP 1002002	A1	20000524	EP 1998-924928	19980528
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 2002511897	T2	20020416	JP 1999-504437	19980528
US 6133325	A	20001017	US 1999-252147	19990218

RUSSEL 09/807,980

PRIORITY APPLN. INFO.:

US 1997-877649 A 19970617

WO 1998-US10814 W 19980528

AB The present invention relates to improved methods for making and using bioadhesive, bioresorbable, antiadhesion compns. made of intermacromol. complexes of carboxyl-contg. polysaccharides and polyethers, and to the resulting compns. The polymers are assocd. with each other, and are then either dried or are used as fluids. Bioresorbable, bioadhesive, antiadhesion compns. are useful in surgery to prevent the formation of post-surgical adhesions. The compns. are designed to breakdown in vivo, and thus be removed from the body. Membranes are inserted during surgery either dry or optionally after conditioning in aq. solns. The antiadhesion, bioadhesive, bioresorptive, antithrombogenic and phys. properties of such membranes can be varied as needed by carefully adjusting the pH of the polymer casting solns., polysaccharide compn., the polyether compn., or by conditioning the membranes prior to surgical use. Bi- or multi-layered membranes can be made and used to provide further control over the phys. and biol. properties of antiadhesion membranes. Antiadhesion compns. may also be used to deliver drugs to the surgical site and release them locally.

IT 1398-61-4, Chitin 9000-69-5, Pectin 9004-32-4, Sodium CMC 9004-42-6, Carboxyethyl cellulose 9004-61-9, Hyaluronic acid 9005-25-8, Starch, biological studies 9005-32-7, Alginic acid 9005-49-6, Heparin, biological studies 9005-79-2, Glycogen, biological studies 9007-28-7, Chondroitin sulfate 9044-05-7, Carboxymethyl dextran 9050-30-0, Heparan sulfate 25322-68-3, Polyethylene oxide 83512-85-0, Carboxymethyl chitosan  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(bioresorbable adhesives contg. **carboxypolysaccharide** -polyether intermacromol. complexes)

RN 1398-61-4 HCAPLUS

CN Chitin (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9000-69-5 HCAPLUS

CN Pectin (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9004-32-4 HCAPLUS

CN Cellulose, carboxymethyl ether, sodium salt (8CI, 9CI) (CA INDEX NAME)

CM 1

CRN 9004-34-6

CMF Unspecified


CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1

CMF C2 H4 O3



RN 9004-42-6 HCAPLUS

RUSSEL 09/807,980

CN Cellulose, 2-carboxyethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 9004-34-6

CMF Unspecified


CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 503-66-2

CMF C3 H6 O3



RN 9004-61-9 HCAPLUS

CN Hyaluronic acid (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9005-25-8 HCAPLUS

CN Starch (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9005-32-7 HCAPLUS

CN Alginic acid (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9005-49-6 HCAPLUS

CN Heparin (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9005-79-2 HCAPLUS

CN Glycogen (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9007-28-7 HCAPLUS

CN Chondroitin, hydrogen sulfite (9CI) (CA INDEX NAME)

CM 1

CRN 9007-27-6

CMF Unspecified


CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 7664-93-9

CMF H2 O4 S



RN 9044-05-7 HCAPLUS  
CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)


CM 1

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
CMF C2 H4 O3



RN 9050-30-0 HCAPLUS  
CN Heparan, sulfate (9CI) (CA INDEX NAME)


CM 1

CRN 70226-44-7  
CMF Unspecified  
CCI MAN


\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 7664-93-9  
CMF H2 O4 S



RN 25322-68-3 HCAPLUS  
CN Poly(oxy-1,2-ethanediyl), .alpha.-hydro-.omega.-hydroxy- (9CI) (CA INDEX NAME)



RN 83512-85-0 HCAPLUS  
CN Chitosan, N-(carboxymethyl) (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

IT 7664-41-7, Ammonia, uses  
RL: NUU (Other use, unclassified); USES (Uses)  
(membrane conditioning with; bioresorbable adhesives contg.  
carboxypolysaccharide-polyether intermacromol. complexes)

RN 7664-41-7 HCAPLUS  
CN Ammonia (8CI, 9CI) (CA INDEX NAME)

NH3

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L47 ANSWER 7 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1998:665874 HCAPLUS

DOCUMENT NUMBER: 130:4084

TITLE: Preparation of **polysaccharide-peptide** or  
amino acid-linked camptothecin **conjugates** as  
antitumor agents

INVENTOR(S): Tsujihara, Kenji; Kawaguchi, Takayuki; Okuno, Akira;  
Yano, Toshiaki

PATENT ASSIGNEE(S): Tanabe Seiyaku Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 44 pp.

CODEN: JKXXAF

DOCUMENT TYPE: Patent



LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 10273488	A2	19981013	JP 1998-16763	19980129
PRIORITY APPLN. INFO.:			JP 1997-17280	19970131
OTHER SOURCE(S):	MARPAT 130:4084			

GI



AB The title compds., which are camptothecin derives. [I; R1 =

(un)substituted lower alkyl; X1 = NHR2, OH; wherein R2 = H, lower alkyl; Alk = linear or branched alkylene optionally interrupted by O] linked to carboxy-contg. polysaccharide through a peptide or amino acid, are prepd. These compds. are reduced in toxicity and markedly enhanced in antitumor potency. Claimed is a pharmaceutical compn. contg. I as the active ingredient for treatment of cancers of lung, uterus, ovary, breast, digestive organs (large intestine, stomach, or pancreas), liver, kidney, prostate gland, and neck, malignant lymphoma, and leukemia. Thus, N-peptidyl-10-(3-aminopropoxy)-(20S)-camptothecin deriv. (II; R = H) (prepn. given) was condensed with carboxymethyl dextran sodium salt using 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride in H2O to give the title compd. II (R = carboxymethyl dextran sodium salt residue), which at 60 mg/kg (single dosage) in vivo inhibited 100% the proliferation of human breast cancer MX-1 cell in mice within 26 days after the drug administration.

IT 39422-83-8DP, Carboxymethyl dextran sodium salt, conjugates with peptide-linked camptothecin derivs.  
 53571-87-2DP, Carboxymethyl pullulan, conjugates with peptide-linked camptothecin derivs., sodium salt 187793-65-3P  
 187793-71-1P 187794-13-4P 187794-21-4P  
 187794-24-7P 187794-27-0P 187794-30-5P  
 187794-33-8P 187794-36-1P 187803-18-5DP, bound to carboxymethyl dextran sodium salt  
 187803-20-9DP, bound to carboxymethyl dextran sodium salt 187803-21-0DP, bound to carboxymethyl dextran sodium salt 187803-22-1DP, bound to carboxymethyl dextran sodium salt 187803-23-2DP  
 , bound to carboxymethyl dextran sodium salt 187803-26-5DP, bound to carboxymethyl dextran sodium salt 187803-27-6DP, bound to carboxymethyl dextran sodium salt 187803-28-7DP, bound to carboxymethyl dextran sodium salt 187803-29-8DP  
 , bound to carboxymethyl dextran sodium salt 187803-30-1DP, bound to carboxymethyl dextran sodium salt 187803-31-2DP, bound to carboxymethyl dextran sodium salt 187803-32-3DP, bound to carboxymethyl dextran sodium salt 187803-33-4DP  
 , bound to carboxymethyl dextran sodium salt 187803-34-5DP, bound to carboxymethyl dextran sodium salt 187803-35-6DP, bound to carboxymethyl dextran sodium salt 215591-97-2DP, bound to carboxymethyl dextran sodium salt 215591-98-3DP  
 , bound to carboxymethyl dextran sodium salt 215592-03-3P 215592-06-6P 215592-09-9P  
 215592-15-7P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (prepn. of polysaccharide-peptide or amino acid-linked camptothecin conjugates as antitumor agents)

RN 39422-83-8 HCAPLUS  
 CN Dextran, carboxymethyl ether, sodium salt (9CI) (CA INDEX NAME)


CM 1

CRN 9004-54-0  
 CMF Unspecified  
 CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
CMF C2 H4 O3



RN 53571-87-2 HCAPLUS  
CN Pullulan, carboxymethyl ether (9CI) (CA INDEX NAME)


CM 1

CRN 9057-02-7  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE


CM 2

CRN 79-14-1  
CMF C2 H4 O3



RN 187793-65-3 HCAPLUS  
CN Benzenepropanamide, .alpha.-amino-N-[3-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-4-hydroxy-, monohydrochloride, (.alpha.S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● HCl

RN 187793-71-1 HCAPLUS  
CN Propanamide, 2-amino-N-[3-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-

RUSSEL 09/807,980

yl]oxy]propyl]-3-hydroxy-, monohydrochloride, (2S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● HCl

RN 187794-13-4 HCAPLUS

CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione,  
4,11-diethyl-4-hydroxy-9-[2-(2-hydroxyethoxy)ethoxy]-, (4S)- (9CI) (CA  
INDEX NAME)

Absolute stereochemistry.




RN 187794-21-4 HCAPLUS

CN L-Alanine, 3-[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-  
1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl ester,  
monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.






● HCl

RN 187794-24-7 HCAPLUS

CN L-Alanine, 2-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethyl ester, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● HCl

RN 187794-27-0 HCAPLUS

CN L-Alanine, 2-[2-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethoxy]ethyl ester, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.



● HCl

RN 187794-30-5 HCAPLUS  
 CN L-Proline, 3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl ester, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.



● HCl

RN 187794-33-8 HCAPLUS  
 CN L-Proline, 2-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethyl ester, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.



● HCl

RN 187794-36-1 HCAPLUS  
 CN L-Proline, 2-[2-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethoxy]ethyl ester, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.





● HCl

RN 187803-18-5 HCAPLUS  
 CN Glycinamide, glycyglycyl-L-phenylalanyl-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A






RN 187803-20-9 HCAPLUS

CN Glycinamide, L-phenylalanyl-N-[3-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 187803-21-0 HCAPLUS

CN Glycinamide, glycylglycyl-L-phenylalanyl-N-[2-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




PAGE 1-B




RN 187803-22-1 HCAPLUS  
 CN Glycinamide, glycyglycyl-L-phenylalanyl-N-[3-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-N-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A




PAGE 1-B




RN 187803-23-2 HCAPLUS  
 CN Glycinamide, glycyglycylglycyl-N-[3-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





RN 187803-26-5 HCAPLUS

CN Glycinamide, glycylglycyl-L-phenylalanyl-N-[2-[2-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethoxy]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A







RN 187803-27-6 HCAPLUS

CN Glycine, glycyglycyl-L-phenylalanyl-, 3-[[4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187803-28-7 HCAPLUS

CN Acetamide, 2-amino-N-[3-[[4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 187803-29-8 HCAPLUS


CN Glycinamide, glycylglycyl-L-phenylalanyl-N-[5-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]pentyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B





RN 187803-30-1 HCAPLUS

CN Glycinamide, glycylglycyl-L-phenylalanyl-N-[4-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]butyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.






RN 187803-31-2 HCAPLUS

CN Glycinamide, glycyL-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187803-32-3 HCAPLUS

CN Glycinamide, D-phenylalanyl-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 187803-33-4 HCAPLUS


CN Glycinamide, glycyglycyl-D-phenylalanyl-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A





PAGE 1-B



RN 187803-34-5. HCAPLUS

CN Glycinamide, glycyglycyl-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 187803-35-6 HCAPLUS

CN Glycinamide, glycylglycylglycylglycyl-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.






RN 215591-97-2 HCAPLUS

CN Butanamide, 4-amino-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 215591-98-3 HCAPLUS

CN Butanamide, 4-[(4-amino-1-oxobutyl)amino]-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.






RN 215592-03-3 HCAPLUS

CN L-Aspartic acid, 4-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl] 1-ethyl ester, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● HCl

RN 215592-06-6 HCAPLUS

CN L-Aspartic acid, 4-[2-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethyl] 1-ethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 215592-09-9 HCAPLUS

CN L-Aspartic acid, 4-[2-[2-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethoxy]ethyl] 1-ethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A




PAGE 1-B

—OEt


RN 215592-15-7 HCAPLUS  
 CN Glycinamide, glycyl-L-phenylalanylglycyl-N-[3-[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-N-methyl-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.


PAGE 1-A




● HCl




IT 79-11-8, Chloroacetic acid, reactions 98-59-9, Tosyl chloride 156-87-6, 3-Aminopropanol 627-30-5, 3-Chloropropanol 1826-67-1, Vinylmagnesium bromide 3978-80-1 9004-54-0, Dextran, reactions 9057-02-7, Pullulan 15761-38-3 17302-47-5 18162-48-6, tert-Butyldimethylsilyl chloride 24424-99-5, Di-tert-butyl dicarbonate 28782-81-2 42454-06-8, 5-Hydroxy-2-nitrobenzaldehyde 110351-94-5  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (prepn. of **polysaccharide-peptide** or amino acid-linked camptothecin **conjugates** as antitumor agents)  
 RN 79-11-8 HCAPLUS  
 CN Acetic acid, chloro- (8CI, 9CI) (CA INDEX NAME)




RN 98-59-9 HCAPLUS  
 CN Benzenesulfonyl chloride, 4-methyl- (9CI) (CA INDEX NAME)



RN 156-87-6 HCAPLUS  
 CN 1-Propanol, 3-amino- (8CI, 9CI) (CA INDEX NAME)




RN 627-30-5 HCAPLUS  
 CN 1-Propanol, 3-chloro- (6CI, 7CI, 8CI, 9CI) (CA INDEX NAME)



RN 1826-67-1 HCAPLUS


CN Magnesium, bromoethenyl- (9CI) (CA INDEX NAME)



RN 3978-80-1 HCAPLUS

CN L-Tyrosine, N-[(1,1-dimethylethoxy)carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



RN 9004-54-0 HCAPLUS

CN Dextran (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9057-02-7 HCAPLUS


CN Pullulan (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 15761-38-3 HCAPLUS


CN L-Alanine, N-[(1,1-dimethylethoxy)carbonyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 17302-47-5 HCAPLUS


CN Phenol, 3-(dimethoxymethyl)-4-nitro- (9CI) (CA INDEX NAME)




RN 18162-48-6 HCAPLUS

CN Silane, chloro(1,1-dimethylethyl)dimethyl- (9CI) (CA INDEX NAME)






RN 24424-99-5 HCAPLUS  
CN Dicarmonic acid, bis(1,1-dimethylethyl) ester (9CI) (CA INDEX NAME)




RN 28782-81-2 HCAPLUS  
CN Glycine, N-[(1,1-dimethylethoxy)carbonyl]-L-phenylalanyl-L-alanyl- (9CI)  
(CA INDEX NAME)

Absolute stereochemistry.




RN 42454-06-8 HCAPLUS  
CN Benzaldehyde, 5-hydroxy-2-nitro- (7CI, 9CI) (CA INDEX NAME)



RN 110351-94-5 HCAPLUS  
CN 1H-Pyrano[3,4-f]indolizine-3,6,10(4H)-trione, 4-ethyl-7,8-dihydro-4-hydroxy-, (4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



IT 39422-83-8P, Carboxymethyl dextran sodium salt  
53571-87-2DP, Carboxymethyl pullulan, sodium salt  
58885-58-8P 80909-96-2P 187793-42-6P  
187793-43-7P 187793-44-8P 187793-46-0P  
187793-48-2P 187793-52-8P 187793-56-2P

187793-58-4P 187793-60-8P 187793-62-0P  
 187793-67-5P 187793-69-7P 187793-76-6P  
 187793-80-2P 187793-82-4P 187793-84-6P  
 187793-86-8P 187794-01-0P 187794-03-2P  
 187794-05-4P 187794-07-6P 187794-09-8P  
 187794-11-2P 187794-17-8P 187794-19-0P  
 187794-20-3P 187794-22-5P 187794-23-6P  
 187794-25-8P 187794-26-9P 187794-28-1P  
 187794-29-2P 187794-31-6P 187794-32-7P  
 187794-34-9P 187794-35-0P 187794-47-4P  
 187794-50-9P 187794-55-4P 187794-58-7P  
 187794-60-1P 187794-66-7P 187794-68-9P  
 187794-70-3P 187794-72-5P 187794-74-7P  
 187803-36-7P 187803-37-8P 205647-87-6P  
 215591-99-4P 215592-00-0P 215592-01-1P  
 215592-02-2P 215592-04-4P 215592-05-5P  
 215592-07-7P 215592-08-8P 215592-10-2P  
 215592-11-3P 215592-12-4P 215592-13-5P  
 215592-14-6P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT  
 (Reactant or reagent)

(prepn. of **polysaccharide**-peptide or amino acid-linked  
 camptothecin **conjugates** as antitumor agents)

RN 39422-83-8 HCAPLUS

CN Dextran, carboxymethyl ether, sodium salt (9CI) (CA INDEX NAME)

CM 1

CRN 9004-54-0

CMF Unspecified


CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1

CMF C2 H4 O3



RN 53571-87-2 HCAPLUS

CN Pullulan, carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 9057-02-7

CMF Unspecified


CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2


CRN 79-14-1

CMF C2 H4 O3




RN 58885-58-8 HCAPLUS

CN Carbamic acid, (3-hydroxypropyl)-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)




RN 80909-96-2 HCAPLUS

CN Carbamic acid, [3-[[4-methylphenyl)sulfonyl]oxy]propyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)




RN 187793-42-6 HCAPLUS

CN Benzenemethanol, .alpha.-ethenyl-5-hydroxy-2-nitro- (9CI) (CA INDEX NAME)




RN 187793-43-7 HCAPLUS

CN Carbamic acid, [3-[3-(1-hydroxy-2-propenyl)-4-nitrophenoxy]propyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)



RN 187793-44-8 HCAPLUS


CN Carbamic acid, [3-[4-nitro-3-(1-oxo-2-propenyl)phenoxy]propyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)



RN 187793-46-0 HCAPLUS

CN Carbamic acid, [3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187793-48-2 HCAPLUS

CN Acetamide, N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187793-52-8 HCAPLUS

CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione, 9-(3-aminopropoxy)-4,11-diethyl-4-hydroxy-, monohydrochloride, (4S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



● HCl

RN 187793-56-2 HCAPLUS  
 CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione,  
 9-[(5-aminopentyl)oxy]-4,11-diethyl-4-hydroxy-, monohydrochloride, (4S)-  
 (9CI) (CA INDEX NAME)


Absolute stereochemistry.



● HCl

RN 187793-58-4 HCAPLUS  
 CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione,  
 9-[2-(2-aminoethoxy)ethoxy]-4,11-diethyl-4-hydroxy-, monohydrochloride,  
 (4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



● HCl


RN 187793-60-8 HCAPLUS  
 CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione,

searched by Susan Hanley 305-4053

RUSSEL 09/807,980

4,11-diethyl-4-hydroxy-9-[3-(methylamino)propoxy]-, monohydrochloride,  
(4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● HCl

RN 187793-62-0 HCAPLUS

CN Carbamic acid, [(1S)-2-[[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]amino]-1-[(4-hydroxyphenyl)methyl]-2-oxoethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187793-67-5 HCAPLUS

CN Acetamide, 2-amino-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● HCl

RN 187793-69-7 HCAPLUS

CN Carbamic acid, [(1S)-2-[[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]amino]-1-(hydroxymethyl)-2-oxoethyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187793-76-6 HCAPLUS

CN Glycinamide, L-phenylalanyl-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● HCl

RN 187793-80-2 HCAPLUS

CN Glycinamide, glycyglycyl-L-phenylalanyl-N-[3-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.

PAGE 1-A



● HCl

PAGE 1-B




RN 187793-82-4 HCAPLUS

CN Glycinamide, glycyglycyl-L-phenylalanyl-N-[5-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]pentyl]-, monohydrochloride (9CI) (CA INDEX NAME)




Absolute stereochemistry.

PAGE 1-A



● HCl

PAGE 1-B




RN 187793-84-6 HCAPLUS

CN Glycinamide, glycyglycyl-L-phenylalanyl-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-N-methyl-, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.

PAGE 1-A



● HCl


PAGE 1-B



RN 187793-86-8 HCAPLUS  
 CN Glycinamide, glycyglycyl-L-phenylalanyl-N-[2-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethyl]-, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.

PAGE 1-A




PAGE 1-B

● HCl




RN 187794-01-0 HCAPLUS  
 CN 1-Propanol, 3-[3-(dimethoxymethyl)-4-nitrophenoxy]- (9CI) (CA INDEX NAME)




RN 187794-03-2 HCAPLUS

CN Benzaldehyde, 5-[3-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]propoxy]-2-nitro- (9CI) (CA INDEX NAME)




RN 187794-05-4 HCAPLUS

CN 2-Propen-1-one, 1-[5-[3-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]propoxy]-2-nitrophenyl]- (9CI) (CA INDEX NAME)



RN 187794-07-6 HCAPLUS


CN Benzenemethanol, 5-[3-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]propoxy]-.alpha.-ethenyl-2-nitro- (9CI) (CA INDEX NAME)



RN 187794-09-8 HCAPLUS


CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione, 4,11-diethyl-4-hydroxy-9-(3-hydroxypropoxy)-, (4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 187794-11-2 HCAPLUS

CN 2-Propen-1-one, 1-[5-[2-[2-[(1,1-dimethylethyl)dimethylsilyl]oxy]ethoxy]ethoxy]-2-nitrophenyl]- (9CI) (CA INDEX NAME)



RN 187794-17-8 HCAPLUS


CN 2-Propen-1-one, 1-[5-(3-hydroxypropoxy)-2-nitrophenyl]- (9CI) (CA INDEX NAME)



RN 187794-19-0 HCAPLUS

CN L-Alanine, N-[(1,1-dimethylethoxy)carbonyl]-, 3-[4-nitro-3-(1-oxo-2-propenyl)phenoxy]propyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187794-20-3 HCAPLUS

CN L-Alanine, N-[(1,1-dimethylethoxy)carbonyl]-, 3-[[4S]-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187794-22-5 HCAPLUS

CN L-Alanine, N-[(1,1-dimethylethoxy)carbonyl]-, 2-[4-nitro-3-(1-oxo-2-propenyl)phenoxy]ethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187794-23-6 HCAPLUS

CN L-Alanine, N-[(1,1-dimethylethoxy)carbonyl]-, 2-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187794-25-8 HCAPLUS

CN L-Alanine, N-[(1,1-dimethylethoxy)carbonyl]-, 2-[2-[4-nitro-3-(1-oxo-2-propenyl)phenoxy]ethoxy]ethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 187794-26-9 HCAPLUS

CN L-Alanine, N-[(1,1-dimethylethoxy)carbonyl]-, 2-[2-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethoxy]ethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A




PAGE 1-B

—OBu-t

RN 187794-28-1 HCAPLUS

CN 1,2-Pyrrolidinedicarboxylic acid, 1-(1,1-dimethylethyl) 2-[3-[4-nitro-3-(1-oxo-2-propenyl)phenoxy]propyl] ester, (2S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187794-29-2 HCAPLUS

CN 1,2-Pyrrolidinedicarboxylic acid, 2-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl] 1-(1,1-dimethylethyl) ester, (2S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187794-31-6 HCAPLUS

CN 1,2-Pyrrolidinedicarboxylic acid, 1-(1,1-dimethylethyl) 2-[2-[4-nitro-3-(1-oxo-2-propenyl)phenoxy]ethyl] ester, (2S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187794-32-7 HCAPLUS

CN 1,2-Pyrrolidinedicarboxylic acid, 2-[2-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethyl] 1-(1,1-dimethylethyl) ester, (2S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 187794-34-9 HCAPLUS

CN 1,2-Pyrrolidinedicarboxylic acid, 1-(1,1-dimethylethyl)  
2-[2-[2-[4-nitro-3-(1-oxo-2-propenyl)phenoxy]ethoxy]ethyl] ester, (2S)-  
(9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 187794-35-0 HCAPLUS

CN 1,2-Pyrrolidinedicarboxylic acid, 2-[2-[2-[[4S]-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethoxy]ethyl] 1-(1,1-dimethylethyl) ester, (2S)- (9CI)  
(CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A






PAGE 1-B

—OBU-t


RN 187794-47-4 HCAPLUS  
 CN Glycinamide, glycyglycyglycyl-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A




PAGE 1-B



RN 187794-50-9 HCAPLUS  
 CN Glycinamide, glycyglycyl-L-phenylalanyl-N-[2-[2-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethoxy]ethyl]-, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.

PAGE 1-A



● HCl

PAGE 1-B




RN 187794-55-4 HCAPLUS


CN Glycine, glycyglycyl-L-phenylalanyl-, 3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl ester, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A




● HCl



RN 187794-58-7 HCAPLUS  
 CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione,  
 9-(4-aminobutoxy)-4,11-diethyl-4-hydroxy-, monohydrochloride, (4S)- (9CI)  
 (CA INDEX NAME)


Absolute stereochemistry.




● HCl

RN 187794-60-1 HCAPLUS  
 CN Glycinamide, glycyglycyl-L-phenylalanyl-N-[4-[[[(4S)-4,11-diethyl-  
 3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,  
 2-b]quinolin-9-yl]oxy]butyl]-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● HCl



RN 187794-66-7 HCAPLUS

CN Glycinamide, glycyl-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 187794-70-3 HCAPLUS

CN Glycinamide, glycyglycyl-N-[3-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.

PAGE 1-A



● HCl



PAGE 1-B



RN 187794-72-5 HCAPLUS

CN Glycinamide, glycyglycyglycyglycyl-N-[3-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 187794-74-7 HCAPLUS

CN Glycinamide, glycylglycyl-D-phenylalanyl-N-[3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.





RN 187803-36-7 HCAPLUS


CN 2-Propen-1-one, 1-[5-[2-[[[(1,1-dimethylethyl)dimethylsilyl]oxy]ethoxy]-2-nitrophenyl]- (9CI) (CA INDEX NAME)



RN 187803-37-8 HCAPLUS


CN 1H-Pyrano[3',4':6,7]indolizino[1,2-b]quinoline-3,14(4H,12H)-dione, 4,11-diethyl-4-hydroxy-9-(2-hydroxyethoxy)-, (4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 205647-87-6 HCAPLUS

CN Carbamic acid, [3-[4-amino-3-(1-oxopropyl)phenoxy]propyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)




RN 215591-99-4 HCAPLUS

CN Glycinamide, N-[(1,1-dimethylethoxy)carbonyl]-L-phenylalanyl-N-[3-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



● HCl

PAGE 1-B


—OBu-t

RN 215592-00-0 HCAPLUS


CN Glycinamide, N-[(1,1-dimethylethoxy)carbonyl]glycylglycyl-L-phenylalanyl-N-[3-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.






● HCl



RN 215592-01-1 HCAPLUS

CN Butanamide, 4-amino-N-[3-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● HCl

RN 215592-02-2 HCAPLUS

CN Butanamide, 4-[(4-amino-1-oxobutyl)amino]-N-[3-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.

PAGE 1-A



● HCl


PAGE 1-B



RN 215592-04-4 HCAPLUS

CN L-Aspartic acid, N-[(1,1-dimethylethoxy)carbonyl]-, 1-ethyl  
4-[3-[4-nitro-3-(1-oxo-2-propenyl)phenoxy]propyl] ester (9CI) (CA INDEX  
NAME)


Absolute stereochemistry.



RN 215592-05-5 HCAPLUS

CN L-Aspartic acid, N-[(1,1-dimethylethoxy)carbonyl]-, 4-[3-[[[4S]-4,11-  
diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-  
pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl] 1-ethyl ester  
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 215592-07-7 HCAPLUS

CN L-Aspartic acid, N-[(1,1-dimethylethoxy)carbonyl]-, 1-ethyl  
4-[2-[4-nitro-3-(1-oxo-2-propenyl)phenoxy]ethyl] ester (9CI) (CA INDEX  
NAME)


Absolute stereochemistry.



RN 215592-08-8 HCAPLUS

CN L-Aspartic acid, N-[(1,1-dimethylethoxy)carbonyl]-, 4-[2-[[ (4S)-4,11-  
diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-  
pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethyl] 1-ethyl ester  
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 215592-10-2 HCAPLUS

CN L-Aspartic acid, N-[(1,1-dimethylethoxy)carbonyl]-, 1-ethyl  
4-[2-[2-[4-nitro-3-(1-oxo-2-propenyl)phenoxy]ethoxy]ethyl] ester (9CI)  
(CA INDEX NAME)

Absolute stereochemistry.




RN 215592-11-3 HCAPLUS

CN L-Aspartic acid, N-[(1,1-dimethylethoxy)carbonyl]-, 4-[2-[2-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethoxy]ethyl] 1-ethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B


—OBu-t

—OEt


RN 215592-12-4 HCAPLUS

CN Glycinamide, N-[(1,1-dimethylethoxy)carbonyl]glycylglycylglycyl-N-[3-[[ (4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl]-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● HCl



RN 215592-13-5 HCAPLUS


CN Glycinamide, N-[(1,1-dimethylethoxy)carbonyl]glycylglycyl-L-phenylalanyl-N-[2-[2-[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]ethoxy]ethyl]-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.



● HCl

PAGE 1-B




RN 215592-14-6 HCAPLUS

CN Glycine, N-[(1,1-dimethylethoxy)carbonyl]glycylglycyl-L-phenylalanyl-,  
3-[[[(4S)-4,11-diethyl-3,4,12,14-tetrahydro-4-hydroxy-3,14-dioxo-1H-  
pyrano[3',4':6,7]indolizino[1,2-b]quinolin-9-yl]oxy]propyl ester,  
monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.

PAGE 1-A



● HCl

PAGE 1-B



L47 ANSWER 8 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1998:127052 HCAPLUS

DOCUMENT NUMBER: 128:172026

searched by Susan Hanley 305-4053

TITLE: Extending Insulin Action in Vivo by  
**Conjugation to Carboxymethyl  
Dextran**

AUTHOR(S): Baudys, Miroslav; Letourneur, Didier; Liu, Feng; Mix,  
Don; Jozefonvicz, Jacqueline; Kim, Sung Wan

CORPORATE SOURCE: Department of Pharmaceutics and Pharmaceutical  
Chemistry/Center for Controlled Chemical Delivery,  
University of Utah, Salt Lake City, UT, 84112, USA

SOURCE: Bioconjugate Chemistry (1998), 9(2), 176-183  
CODEN: BCCHE; ISSN: 1043-1802

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The biochem. and pharmacol. properties of bioactive peptides and proteins can be altered by **conjugation** with polymers. This report describes site-specific attachment of insulin to activated carboxyl groups of **carboxymethyl dextran (CMD, MW = 51 000)** through the GlyAl insulin amino group. On av., three or four insulin mols. were grafted to a **CMD** linear chain. Coupled insulin mols. were properly folded, and the bioactivity of **conjugated** insulin in the blood **glucose** depression assay was 9.6 IU/mg, which was only 2.6 times less than that for native insulin. The cell growth study indicated that the **CMD-insulin conjugate** was as mitogenic as insulin on vascular smooth muscle cells, whereas the starting **CMD** polymer was not. The insulin receptor binding const. of the **conjugate** (3.6 .times. 10<sup>9</sup> M<sup>-1</sup>) compared well with that of native insulin (7.6 .times. 10<sup>9</sup> M<sup>-1</sup>), indicating that the **CMD** chain does not present any major constraints to binding. Plasma clearance of **CMD-insulin** obeyed a two-compartment pharmacokinetic (PK) model with a **CMD-insulin conjugate** plasma elimination half-life of 114.1 min, which was significantly longer than that of sol. Zn-insulin (12.4 min). In contrast, pharmacodynamic (PD) profiles (blood **glucose** lowering effects) after i.v. (i.v.) administration of the **conjugate** or insulin in rats were not different. S.c. (s.c.) administration of the **conjugate** resulted in a significantly prolonged plasma profile with a noncompartmental PK parameter mean residence time (MRT) of 103.5 min which was significantly longer than that of sol. Zn-insulin (40.5 min). This was reflected in the protracted PD effect of s.c. administered **conjugate** with time needed to reach min. **glucose** concn. Tnadir of 95.7 min, which was significantly longer than that of insulin (62 min). We conclude that the **conjugation** of insulin to **CMD** leads to a bioactive **conjugate** with a delayed s.c. PD profile showing prolonged response, resembling intermediate acting insulin preps.

IT 9004-10-8DP, Insulin, **conjugates** with **carboxymethyl dextran**, biological studies  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)  
(extending insulin action in vivo by **conjugation** to **carboxymethyl dextran**)

RN 9004-10-8 HCAPLUS  
CN Insulin (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

IT 9044-05-7DP, **Carboxymethyl dextran**, **conjugates** with insulin  
RL: BPR (Biological process); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study);

PREP (Preparation); PROC (Process); USES (Uses)  
(extending insulin action in vivo by **conjugation** to  
**carboxymethyl dextran**)

RN 9044-05-7 HCAPLUS

CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 9004-54-0

CMF Unspecified


CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1

CMF C2 H4 O3



L47 ANSWER 9 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1997:777835 HCAPLUS

DOCUMENT NUMBER: 128:97464

TITLE: Design of macromolecular biological response modifier  
by immobilizing of D-glucose analog of muramyl  
dipeptide on carboxymethyl-dextran having mannose  
branches

AUTHOR(S): Murata, J.; Nagae, H.; Ohya, Y.; Ouchi, T.

CORPORATE SOURCE: Dep. Applied Chem., Faculty Eng., Kansai Univ., Suita,  
564, Japan

SOURCE: Journal of Biomaterials Science, Polymer Edition  
(1997), 8(12), 931-946

CODEN: JBSEEA; ISSN: 0920-5063

PUBLISHER: VSP BV

DOCUMENT TYPE: Journal

LANGUAGE: English


AB It is well known that muramyl dipeptide is a min. required structure of  
bacterial peptidoglycan responsible for immunoadjuvant activity. Since  
mannose receptors exists on the surface of macrophages, polymers with  
branched mannose residues are expected to target moieties to macrophages.  
To achieve an efficient delivery of D-**glucose** analog of muramyl  
dipeptide (GADP) via receptor-mediated endocytosis by mannose receptors on  
the surface of macrophages, GADP/**carboxymethyl-dextran**  
(CM-Dex)/Man **conjugate** was synthesized. Moreover, to study the  
effect of the introduction of mannose residues, we also synthesized  
GADP/CM-glucomannan (CM-GM) and GADP/CM-Dex **conjugates**. The  
immunol. enhancement activities of their **conjugates** were  
evaluated by measurements of **glucose** consumption and  
.beta.-D-glucuronidase activity from macrophage-like cells. The  
GADP/CM-Dex/Man and GADP/CM-GM **conjugates** showed higher immunol.  
enhancement activity than the GADP/CM-Dex **conjugate**. The  
immunol. enhancement activity of GADP/CM-Dex/Man and GADP/CM-GM  
**conjugates** was decreased to the same level of immunol enhancement  
activity of GADP/CM-Dex **conjugate** under the presence of excess  
mannose. These results suggested that the introduction of mannose



residues into GADP/CM-Dex **conjugate** could increase the affinity against macrophage and the immunol. enhancement activity of GADP/CM-Dex **conjugate** itself.


IT **146916-64-5DP**, reaction products with CM-dextran or CM-glucomannan  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (design of macromol. biol. response modifier by immobilizing glucose analog of muramyl dipeptide on CM-dextran having mannose branches)  
 RN 146916-64-5 HCAPLUS  
 CN D-.alpha.-Glutamine, N-[(2R)-2-[6-O-(6-amino-1-oxohexyl)-D-glucos-3-O-yl]-1-oxopropyl]-L-alanyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT **50-99-7**, D-Glucose, reactions **617-04-9**, Methyl .alpha.-D-mannopyranoside **6404-29-1** **9044-05-7D**, Carboxymethyl dextran, reaction products with glucose analog of muramyl dipeptide **9064-52-2D**, Carboxymethyl glucomannan, reaction products with glucose analog of muramyl dipeptide **77987-49-6**, N-Benzoyloxycarbonyl ethanolamine **107947-55-7**  
 RL: RCT (Reactant); RACT (Reactant or reagent)  
 (design of macromol. biol. response modifier by immobilizing glucose analog of muramyl dipeptide on CM-dextran having mannose branches)  
 RN 50-99-7 HCAPLUS  
 CN D-Glucose (8CI, 9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 617-04-9 HCAPLUS  
 CN .alpha.-D-Mannopyranoside, methyl (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 6404-29-1 HCAPLUS

RUSSEL 09/807,980

CN Hexanoic acid, 6-[[ (1,1-dimethylethoxy)carbonyl]amino]- (9CI) (CA INDEX NAME)



RN 9044-05-7 HCAPLUS

CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 9004-54-0

CMF Unspecified


CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1

CMF C2 H4 O3



RN 9064-52-2 HCAPLUS

CN D-Gluco-D-mannan, carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 11078-31-2

CMF Unspecified


CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2


CRN 79-14-1

CMF C2 H4 O3




RN 77987-49-6 HCAPLUS

CN Carbamic acid, (2-hydroxyethyl)-, phenylmethyl ester (9CI) (CA INDEX NAME)




RN 107947-55-7 HCAPLUS  
 CN D-.alpha.-Glutamine, L-alanyl-, (4-nitrophenyl)methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 74112-33-7P 78609-16-2P, .alpha.-D-Mannopyranose, 2,3,4,6-tetrakis-O-(phenylmethyl)- 92470-93-4P, 2,3,4,6-Tetra-O-benzyl-.alpha.-D-mannopyranosyl chloride 140428-88-2DP, reaction products with CM-dextran and glucose analog of muramyl dipeptide 140428-88-2P 146916-64-5P 201145-84-8P 201145-85-9P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)  
 (design of macromol. biol. response modifier by immobilizing glucose analog of muramyl dipeptide on CM-dextran having mannose branches)  
 RN 74112-33-7 HCAPLUS  
 CN .alpha.-D-Glucofuranose, 3-O-(1-carboxyethyl)-1,2:5,6-bis-O-(1-methylethylidene)-, (R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 78609-16-2 HCAPLUS  
 CN .alpha.-D-Mannopyranose, 2,3,4,6-tetrakis-O-(phenylmethyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



RN 92470-93-4 HCAPLUS  
 CN .alpha.-D-Mannopyranosyl chloride, 2,3,4,6-tetrakis-O-(phenylmethyl)-

(9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 140428-88-2 HCAPLUS.

CN .alpha.-D-Mannopyranoside, 2-aminoethyl (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



RN 140428-88-2 HCAPLUS

CN .alpha.-D-Mannopyranoside, 2-aminoethyl (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



RN 146916-64-5 HCAPLUS

CN D-.alpha.-Glutamine, N-[(2R)-2-[6-O-(6-amino-1-oxohexyl)-D-glucos-3-O-yl]-1-oxopropyl]-L-alanyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 201145-84-8 HCAPLUS

CN Carbamic acid, [2-[[2,3,4,6-tetrakis-O-(phenylmethyl)-.alpha.-D-mannopyranosyl]oxy]ethyl]-, phenylmethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 201145-85-9 HCAPLUS

CN D-.alpha.-Glutamine, N-[(2R)-2-[1,2:5,6-bis-O-(1-methylethylidene)-.alpha.-D-glucofuranos-3-O-yl]-1-oxopropyl]-L-alanyl-, (4-nitrophenyl)methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L47 ANSWER 10 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1997:138228 HCAPLUS

DOCUMENT NUMBER: 126:242714

TITLE: Targeted delivery of drugs and proteins to the liver via receptor-mediated endocytosis

AUTHOR(S): Hashida, Mitsuru; Hirabayashi, Hideki; Nishikawa, Makiya; Takakura, Yoshinobu

CORPORATE SOURCE: Department of Drug Delivery Research, Faculty of Pharmaceutical Sciences, Kyoto University, Sakyo-ku, Kyoto, Japan

SOURCE: J. Controlled Release (1997), 46(1,2), 129-137

CODEN: JCREEC; ISSN: 0168-3659

PUBLISHER: Elsevier

DOCUMENT TYPE: Journal

LANGUAGE: English


AB Targeting of drugs and proteins to the liver via the asialoglycoprotein receptor was investigated in mice. **Carboxymethyl-dextran (CMD)**, carboxymethyl-amylose (CMA), and poly-L-glutamic acid (PLGA) were modified with 2-imino-2-methoxyethyl (IME)-thiogalactosides to obtain galactosylated derivs. as carriers of drugs with low-mol. wts. Proteins were targeted to the liver by direct attachment of **galactose** moieties. Pharmacokinetic anal. clearly showed that galactosylated derivs. were taken up by the liver depending on the mol. wt. and configuration of macromols., the no. of **galactose** residues, and the administered dose. Based on the obtained results, we attempted to selectively deliver vitamin K5, which acts as a coagulant in

the liver. Galactosylated PLGA (Gal-PLGA) possessing 18 **galactose** residues was selected as a hepatotropic carrier since it was efficiently accumulated and gradually degraded in the liver after i.v. injection. The attachment of vitamin K5 did not alter the distribution properties of Gal-PLGA, and vitamin K5 was successfully delivered to the liver by the **conjugation**. The anti-hemorrhagic activity of the **conjugate** was assayed after i.v. injection in mice treated with warfarin. Vitamin K5 **conjugated** with Gal-PLGA showed coagulant activity at any periods studied after i.v. injection, while free vitamin K5 only showed the activity at 4 h after administration. These results indicate the usefulness of galactosylated macromols. as hepatotropic carriers of drugs whose site of action is in the liver.

IT **83-70-5D**, Vitamin K5, reaction products with galactosylated poly(glutamic acid) **107-15-3D**, 1,2-Ethanediamine, reaction products with polysaccharides and galactose deriv. **9044-05-7D**, Carboxymethyl dextran, glycosylated **9054-89-1D**, Superoxide dismutase, galactosylated **12768-31-9D**, Carboxymethyl amylose, glycosylated **24991-23-9D**, glycosylated **25513-46-6D**, Poly(L-glutamic acid), glycosylated **75204-21-6D**, reaction products with macromols.  
 RL: BPR (Biological process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)  
 (targeted delivery of drugs and proteins to the liver via receptor-mediated endocytosis)

RN 83-70-5 HCAPLUS

CN 1-Naphthalenol, 4-amino-2-methyl- (9CI) (CA INDEX NAME)



RN 107-15-3 HCAPLUS

CN 1,2-Ethanediamine (9CI) (CA INDEX NAME)



RN 9044-05-7 HCAPLUS

CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 9004-54-0

CMF Unspecified


CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1

CMF C2 H4 O3



RN 9054-89-1 HCAPLUS  
CN Dismutase, superoxide (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 12768-31-9 HCAPLUS  
CN Amylose, carboxymethyl ether (9CI) (CA INDEX NAME)


CM 1

CRN 9005-82-7  
CMF Unspecified  
CCI PMS, MAN


\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
CMF C2 H4 O3



RN 24991-23-9 HCAPLUS  
CN Poly[imino[(1S)-1-(2-carboxyethyl)-2-oxo-1,2-ethanediyl]] (9CI) (CA INDEX NAME)




RN 25513-46-6 HCAPLUS  
CN L-Glutamic acid, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 56-86-0  
CMF C5 H9 N O4  
CDES 5:L


Absolute stereochemistry.



RN 75204-21-6 HCAPLUS  
CN Ethanimidic acid, 2-(.beta.-D-galactopyranosylthio)-, methyl ester (9CI)

(CA INDEX NAME)

Absolute stereochemistry.



L47 ANSWER 11 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1996:524125 HCAPLUS

DOCUMENT NUMBER: 125:219918

TITLE: Role of the **Polysaccharide** Content and Net Charge on the Emulsifying Properties of .beta.-Lactoglobulin-Carboxymethyldextran **Conjugates**

AUTHOR(S): Nagasawa, Koichi; Ohgata, Koki; Takahashi, Koji; Hattori, Makoto

CORPORATE SOURCE: Faculty of Agriculture, Tokyo University of Agriculture and Technology, Tokyo, 183, Japan

SOURCE: J. Agric. Food Chem. (1996), 44(9), 2538-2543  
CODEN: JAFCAU; ISSN: 0021-8561

DOCUMENT TYPE: Journal

LANGUAGE: English

AB .beta.-Lactoglobulin (.beta.-LG)-carboxymethyldextran (**CMD**) **conjugates** were prepd. by using water-sol. carbodiimide. Three kinds of **CMD** differing in mol. mass (40, 70, and 162 kDa) were used to investigate the effects of different **CMD** contents and net charge on the functional changes in .beta.-LG. The emulsifying properties of these .beta.-LG-**CMD** **conjugates** were investigated under various conditions by evaluating the stability of oil/water emulsions prepd. with oleic acid and the .beta.-LG-**CMD** **conjugates**. The emulsifying ability of .beta.-LG was greatly improved by **conjugating** with **CMD** in the acidic pH range in the presence of less than 0.5 M NaCl. After heating at 80 .degree.C for 10 min, the emulsifying stability of the .beta.-LG-**CMD** **conjugates** was higher than that of .beta.-LG. It is thought that increasing the **polysaccharide** content and shifting the isoelec. point of .beta.-LG to the acidic side by **conjugating** with **CMD** of a high mol. wt. would be effective in improving the emulsifying properties of .beta.-LG under unfavorable conditions.

IT 7647-14-5, Sodium chloride, biological studies

RL: BAC (Biological activity or effector, except adverse); BIOL (Biological study)

(**polysaccharide** content and net charge effect on emulsifying properties of .beta.-lactoglobulin-carboxymethyldextran **conjugates**)

RN 7647-14-5 HCAPLUS

CN Sodium chloride (NaCl) (9CI) (CA INDEX NAME)

Cl-Na



IT 9044-05-7D, Carboxymethyldextran, **conjugates** with  
.beta.-lactoglobulin  
RL: BSU (Biological study, unclassified); PEP (Physical, engineering or  
chemical process); BIOL (Biological study); PROC (Process)  
(**polysaccharide** content and net charge effect on emulsifying  
properties of .beta.-lactoglobulin-carboxymethyldextran  
**conjugates**)

RN 9044-05-7 HCAPLUS

CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 9004-54-0

CMF Unspecified


CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1

CMF C2 H4 O3



L47 ANSWER 12 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1996:507742 HCAPLUS

DOCUMENT NUMBER: 125:204231

TITLE: Pharmacokinetics and targeted delivery of proteins and  
genes

AUTHOR(S): Hashida, Mitsuru; Mahato, Ram I.; Kawabata, Kenji;  
Miyao, Takenori; Nishikawa, Makiya; Takakura,  
Yoshinobu

CORPORATE SOURCE: Fac. Pharmaceutical Sci., Kyoto Univ., Kyoto, 606-01,  
Japan

SOURCE: J. Controlled Release (1996), 41(1,2, Fifth  
International Symposium on Delivery and Targeting of  
Pesticides, Proteins and Genes, 1995), 91-97  
CODEN: JCREEC; ISSN: 0168-3659

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The effectiveness of various approaches for controlling in vivo  
disposition of proteins and genes is compared based on pharmacokinetic  
anal. The potential of introduction of **galactose** or mannose  
residues aiming at receptor-mediated endocytosis, succinylation to be  
recognized by a scavenger receptor, and cationization for universal  
electrostatic interaction were characterized using model proteins.  
Corresponding to the results, a superior therapeutic effect was shown with  
derivs. of superoxide dismutase against hepatic and renal  
ischemia/reperfusion injury. A similar approach was adopted for plasmid  
DNA and oligonucleotide and their rapid degrdn. in the blood pool and  
preferential uptake by the liver after i.v. injection were characterized  
by pharmacokinetic anal. The effects of incorporation into cationic  
liposomes and **conjugation** with macromols. on their in vivo  
distribution were also elucidated.

IT 9044-05-7DP, Carboxymethyldextran, conjugates with 5'-biotinylated

decathymidylic acid 167497-81-6DP, conjugates with  
**carboxymethyl dextran**

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (pharmacokinetics and targeted delivery of proteins and genes)

RN 9044-05-7 HCAPLUS

CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 9004-54-0

CMF Unspecified


CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1


CMF C2 H4 O3





RN 167497-81-6 HCAPLUS

CN Thymidine, P-deoxy-P-[[4-[[5-(hexahydro-2-oxo-1H-thieno[3,4-d]imidazol-4-yl)-1-oxopentyl]amino]butyl]amino]thymidylyl-(3'.fwdarw.5')-thymidylyl-(3'.fwdarw.5')-thymidylyl-(3'.fwdarw.5')-thymidylyl-(3'.fwdarw.5')-thymidylyl-(3'.fwdarw.5')-thymidylyl-(3'.fwdarw.5')-thymidylyl-(3'.fwdarw.5')-, [3aS-(3a.alpha.,4.beta.,6a.alpha.)]- (9CI) (CA INDEX NAME)

PAGE 1-A







PAGE 3-A




PAGE 3-B



PAGE 4-A



PAGE 4-B



IT 9001-63-2, Lysozyme 9035-81-8, Trypsin inhibitor  
9054-89-1, Superoxide dismutase

searched by Susan Hanley 305-4053

# RUSSEL 09/807,980

RL: BAC (Biological activity or effector, except adverse); THU  
(Therapeutic use); BIOL (Biological study); USES (Uses)  
(pharmacokinetics and targeted delivery of proteins and genes)

RN 9001-63-2 HCAPLUS

CN Lysozyme (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9035-81-8 HCAPLUS

CN Trypsin inhibitor (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9054-89-1 HCAPLUS

CN Dismutase, superoxide (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

L47 ANSWER 13 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1996:478465 HCAPLUS

DOCUMENT NUMBER: 125:204200

TITLE: Synthesis of muramyl dipeptide analog-glucomannan  
conjugate and its stimulation activity against  
macrophage-like cells

AUTHOR(S): Murata, Jun-ichi; Nagae, Hiromu; Ohya, Yuichi; Ouchi,  
Tatsuro

CORPORATE SOURCE: Department of Applied Chemistry, Kansai University,  
Suita, 564, Japan

SOURCE: Carbohydr. Polym. (1996), 29(2), 111-118  
CODEN: CAPOD8; ISSN: 0144-8617

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Since the mannose receptors exist on the surface of macrophages, the  
branched mannose residues of glucomannan are expected to act as targeting  
moieties to macrophages. So, to achieve an efficient delivery of D-  
**glucose** analog of muramyl dipeptide (GADP) via receptor-mediated  
endocytosis by mannose receptors on the surface of macrophages, the  
GADP/carboxymethyl (CM)-glucomannan **conjugate** was synthesized.  
Moreover, to study the relation between the immunol. enhancement activity  
of the **conjugates** and their mannose residues, we synthesized the  
GADP/CM-glucomannan **conjugates** having various degrees of  
substitution of carboxymethyl group in mol% per sugar unit (DCM) and  
GADP/CM-dextran **conjugate** through hybridization of GADP with  
dextran. The immunol. enhancement activities of GADP/CM-glucomannan  
**conjugates** and GADP/CM-dextran **conjugate** were evaluated  
by measurements of the **glucose** consumption, the superoxide anion  
prodn. and the .beta.-D-glucuronidase activity from PMA  
(phorbol-12-myristate-13-acetate)-differentiated HL-60 (human  
promyelocytic leukemia) or U937 (human monoblast leukemia) cells as  
macrophage-like cells.

IT 9044-05-7DP, Carboxymethyl dextran,

**conjugates** with **glucose** analog of muramyl dipeptide

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic  
preparation); THU (Therapeutic use); BIOL (Biological study); PREP  
(Preparation); USES (Uses)

(comparison compd.; synthesis of muramyl dipeptide analog-glucomannan  
**conjugate** and stimulation activity against macrophage-like  
cells)

RN 9044-05-7 HCAPLUS

CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
CMF C2 H4 O3



IT **9064-52-2DP**, Carboxymethyl Glucomannan, **conjugates** with **glucose** analog of muramyl dipeptide **146916-65-6DP**, conjugates with CM glucomannan  
RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(synthesis of muramyl dipeptide analog-glucomannan **conjugate** and stimulation activity against macrophage-like cells)  
RN 9064-52-2 HCAPLUS  
CN D-Gluco-D-mannan, carboxymethyl ether (9CI) (CA INDEX NAME)


CM 1

CRN 11078-31-2  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
CMF C2 H4 O3




RN 146916-65-6 HCAPLUS  
CN D-.alpha.-Glutamine, N2-[N-[2-[6-O-(6-amino-1-oxohexyl)-D-glucos-3-O-yl]-1-oxopropyl]-L-alanyl]-, (R)-, mono(trifluoroacetate) (salt) (9CI) (CA INDEX NAME)

CM 1


CRN 146916-64-5  
CMF C23 H40 N4 O12

Absolute stereochemistry.



CM 2

CRN 76-05-1  
CMF C2 H F3 O2



L47 ANSWER 14 OF 22 HCAPLUS COPYRIGHT 2002 ACS  
 ACCESSION NUMBER: 1995:549521 HCAPLUS  
 DOCUMENT NUMBER: 122:312947  
 TITLE: Method of activating soluble carbohydrate using novel  
 cyanylating reagents for the production of immunogenic  
 constructs  
 INVENTOR(S): Lees, Andrew  
 PATENT ASSIGNEE(S): Henry M. Jackson Foundation for the Advancement of  
 Military Medicine, USA  
 SOURCE: PCT Int. Appl., 48 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 4  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9508348	A1	19950330	WO 1994-US10658	19940921
W: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN				
RW: KE, MW, SD, SZ, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
CA 2171942	AA	19950330	CA 1994-2171942	19940921
AU 9478391	A1	19950410	AU 1994-78391	19940921
AU 678613	B2	19970605		
EP 720485	A1	19960710	EP 1994-929273	19940921
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
JP 09502978	T2	19970325	JP 1994-509897	19940921
PRIORITY APPLN. INFO.: US 1993-124491 A 19930922				
WO 1994-US10658 W 19940921				
AB A process for producing an immunogenic construct comprising activating at				

least one first carbohydrate-contg. moiety with a novel cyanylating reagent and covalently joining said activated first moiety to a second moiety. Immunogenic constructs may be prepd. by this process using either direct **conjugation** of first and second moieties or using indirect **conjugation** with a bifunctional reagent. The first carbohydrate is dextran, Pneumococcal **polysaccharide**, Haemophilus influenzae **polysaccharide**, or a viral or bacterial **polysaccharide**; the second carbohydrate is albumin, pertussis toxoid, tetanus toxoid, malaria-derived peptide p28, antibody, toxoid or toxin; the cyanylating reagent is CDAP, CTEA, and pNPC; and the bifunctional reagent is ethylenediamine, 1,6-hexane diamine, adipic dihydrazide, cystamine, glycine, or lysine. In example, **conjugates** of pertussis toxoid and Pneumococcal type 14 **polysaccharide** or Haemophilus influenzae **polysaccharide**, tetanus toxoid and malaria-derived peptide 28, and monoclonal antibody H.delta.a/1 and aminoethyl **carboxymethyl dextran** were prepd. as vaccines.


IT 51-85-4, Cystamine 56-40-6, Glycine, biological studies  
56-87-1, L-Lysine, biological studies 107-15-3,  
1,2-Ethanediamine, biological studies 124-09-4,  
1,6-Hexanediamine, biological studies 1071-93-8, Adipic  
dihydrazide

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
(Uses)

(bifunctional agent; prepn. of immunogenic carbohydrate constructs with  
cyanylating and bifunctional reagent as vaccine)


RN 51-85-4 HCAPLUS

CN Ethanamine, 2,2'-dithiobis- (9CI) (CA INDEX NAME)



RN 56-40-6 HCAPLUS


CN Glycine (8CI, 9CI) (CA INDEX NAME)



RN 56-87-1 HCAPLUS

CN L-Lysine (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 107-15-3 HCAPLUS

CN 1,2-Ethanediamine (9CI) (CA INDEX NAME)




RN 124-09-4 HCAPLUS




RUSSEL 09/807,980

CN 1,6-Hexanediamine (7CI, 8CI, 9CI) (CA INDEX NAME)



RN 1071-93-8 HCAPLUS

CN Hexanedioic acid, dihydrazide (9CI) (CA INDEX NAME)




IT 1129-38-0, p-Nitrophenylcyanate

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)

(cyanylating agent; prepn. of immunogenic carbohydrate constructs with cyanylating and bifunctional reagent as vaccine)

RN 1129-38-0 HCAPLUS

CN Cyanic acid, 4-nitrophenyl ester (9CI) (CA INDEX NAME)



IT 30684-36-7 59016-56-7, 1-Cyano-4-(dimethylamino)-pyridinium tetrafluoroborate

RL: BUU (Biological use, unclassified); RCT (Reactant); BIOL (Biological study); USES (Uses)

(cyanylating agent; prepn. of immunogenic carbohydrate constructs with cyanylating and bifunctional reagent as vaccine)


RN 30684-36-7 HCAPLUS

CN Ethanaminium, N-cyano-N,N-diethyl-, tetrafluoroborate(1-) (9CI) (CA INDEX NAME)

CM 1

CRN 44795-37-1

CMF C7 H15 N2




CM 2

CRN 14874-70-5

CMF B F4


CCI CCS



RN 59016-56-7 HCAPLUS  
CN Pyridinium, 1-cyano-4-(dimethylamino)-, tetrafluoroborate(1-) (9CI) (CA INDEX NAME)


CM 1

CRN 59016-55-6  
CMF C8 H10 N3





CM 2

CRN 14874-70-5  
CMF B F4  
CCI CCS



IT 163038-69-5DP, conjugates with monoclonal antibody  
163438-78-6DP, conjugates with monoclonal antibody  
RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(prepn. of immunogenic carbohydrate constructs with cyanylating and bifunctional reagent as vaccine)  
RN 163038-69-5 HCAPLUS  
CN L-Lysine, L-cysteinyl-L-asparaginyl-L-isoleucylglycyl-L-lysyl-L-prolyl-L-asparaginyl-L-valyl-L-glutaminyl-L-.alpha.-aspartyl-L-.alpha.-aspartyl-L-glutaminyl-L-asparaginyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 163438-78-6 HCAPLUS  
 CN Dextran, 2-aminoethyl carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 9004-54-0  
 CMF Unspecified  
 CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE


CM 2

CRN 141-43-5  
 CMF C2 H7 N O

H<sub>2</sub>N—CH<sub>2</sub>—CH<sub>2</sub>—OH

CM 3

CRN 79-14-1  
 CMF C2 H4 O3



IT 9004-54-0D, Dextran, conjugates  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (prepn. of immunogenic carbohydrate constructs with cyanylating and  
 bifunctional reagent as vaccine)  
 RN 9004-54-0 HCAPLUS  
 CN Dextran (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

L47 ANSWER 15 OF 22 HCAPLUS COPYRIGHT 2002 ACS  
 ACCESSION NUMBER: 1994:289400 HCAPLUS  
 DOCUMENT NUMBER: 120:289400  
 TITLE: Manipulation of renal disposition of human recombinant  
 superoxide dismutase by chemical modification  
 AUTHOR(S): Mihara, Kiyoshi; Sawai, Kenzo; Takakura, Yoshinobu;  
 Hashida, Mitsuru  
 CORPORATE SOURCE: Fac. Pharm. Sci., Kyoto Univ., Kyoto, 606-01, Japan  
 SOURCE: Biol. Pharm. Bull. (1994), 17(2), 296-301  
 CODEN: BPBLEO; ISSN: 0918-6158  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB The renal disposition characteristics of superoxide dismutase (SOD) and  
 its derivs., including macromol. **conjugates** with  
**polyethylene** glycol and **carboxymethyl-dextran**,  
 cationized deriv., and glycosylated derivs. with **galactose** and  
 mannose, were studied in the isolated perfused rat kidney. Renal  
 disposition processes, such as glomerular filtration, tubular  
 reabsorption, and uptake from the capillary side, were quant. detd. by  
 single-pass indicator diln. expts. under filtering and nonfiltering kidney  
 conditions. Native SOD had a high glomerular filtration rate (40% of that  
 of inulin) and was effectively reabsorbed in the tubules, while no  
 significant uptake was obsd. from capillary side. Macromol.  
**conjugates** showed restricted glomerular filtration due to an  
 increase in mol. size. Cationization of SOD greatly enhanced its assocn.  
 with the tissue, not only from the luminal side but also from the  
 capillary side, based upon electrostatic interaction. Galactosylated and  
 mannosylated SOD showed reduced tubular reabsorption and increased  
 exposure of the luminal surface to the enzyme. In addn., a small but  
 significant uptake of mannosylated SOD from the capillary side was obsd.  
 This uptake was dose-dependent and completely inhibited by mannan,  
 suggesting that mannose receptor-mediated endocytosis existed in the  
 capillary side of the kidney. Thus, the authors can manipulate the renal  
 disposition profiles of SOD by changing its physicochem. or biol.  
 properties through chem. modification.

IT 9054-89-1, Superoxide dismutase  
 RL: BIOL (Biological study)  
 (human recombinant, kidney disposition of, chem. modification  
 manipulation of)  
 RN 9054-89-1 HCAPLUS  
 CN Dismutase, superoxide (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

IT 59-23-4DP, **Galactose**, **conjugates** with  
 superoxide dismutase 79-14-1DP, Glycolic acid, conjugates with


RUSSEL 09/807,980

superoxide dismutase **3458-28-4DP**, Mannose, conjugates with  
superoxide dismutase **9044-05-7DP**, Carboxymethyldextran,  
conjugates with superoxide dismutase **25322-68-3DP**,  
**Polyethylene** glycol, conjugates with superoxide dismutase  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(prepn. and kidney disposition of)

RN 59-23-4 HCAPLUS


CN D-Galactose (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



RN 79-14-1 HCAPLUS


CN Acetic acid, hydroxy- (9CI) (CA INDEX NAME)



RN 3458-28-4 HCAPLUS

CN D-Mannose (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



RN 9044-05-7 HCAPLUS

CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 9004-54-0

CMF Unspecified


CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1


CMF C2 H4 O3



RN 25322-68-3 HCAPLUS

CN Poly(oxy-1,2-ethanediyl), .alpha.-hydro-.omega.-hydroxy- (9CI) (CA INDEX

NAME)



L47 ANSWER 16 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1994:235367 HCAPLUS

DOCUMENT NUMBER: 120:235367

TITLE: Control of the disposition profiles of proteins in the kidney via chemical modification

AUTHOR(S): Takakura, Yoshinobu; Mihara, Kiyoshi; Hashida, Mitsuru

CORPORATE SOURCE: Fac. Pharm. Sci., Kyoto Univ., Kyoto, 606-01, Japan

SOURCE: J. Controlled Release (1994), 28(1-3), 111-19

CODEN: JCREEC; ISSN: 0168-3659

DOCUMENT TYPE: Journal

LANGUAGE: English

AB To construct the strategy to control the renal disposition profiles of protein drugs by chem. modification studies were performed using the perfused rat kidney. Renal disposition processes, i.e., glomerular filtration, tubular reabsorption, and uptake from the vascular side, were quant. detd. by single-pass indicator diln. expts. under filtering and non-filtering conditions. As the first step, the renal disposition characteristics of model protein drugs and macromols. were evaluated. These studies clarified the relationship between physicochem. properties of macromols., such as mol. wt. and elec. charge, and their fate in the kidney in a quant. manner. Based on these findings, an antioxidant enzyme, superoxide dismutase (SOD), selected as a therapeutic agent for various tissue injuries including renal failure mediated by reactive oxygen species, was chem. modified. **Conjugation** with macromols., **polyethylene glycol** and **carboxymethyl dextran**, decreased glomerular filtration of SOD. Cationization enabled the enzyme to distribute to the kidney from the capillary side and to be completely reabsorbed by the tubular epithelium after glomerular filtration based on electrostatic interaction. On the other hand, glycosylation with monosaccharides, **galactose** and mannose, significantly reduced its tubular reabsorption and enhanced its exposure to the luminal surface. Furthermore, the mannosylated deriv. accumulated in the kidney from the vascular side via a mannose-recognition mechanism. Thus, the present study demonstrates that chem. modification is useful for the control of renal disposition characteristics of protein drugs.

IT **9044-05-7D, Carboxymethyl dextran, conjugates** with superoxide dismutase **9054-89-1D**, Superoxide dismutase, conjugates with macromols. **25322-68-3D**, **Polyethylene glycol**, conjugates with superoxide dismutase  
RL: BIOL (Biological study)

(disposition profiles of, in kidney)

RN 9044-05-7 HCAPLUS

CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 9004-54-0


CMF Unspecified

CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2


CRN 79-14-1  
CMF C2 H4 O3



RN 9054-89-1 HCAPLUS  
CN Dismutase, superoxide (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 25322-68-3 HCAPLUS  
CN Poly(oxy-1,2-ethanediyl), .alpha.-hydro-.omega.-hydroxy- (9CI) (CA INDEX NAME)



L47 ANSWER 17 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1994:226756 HCAPLUS

DOCUMENT NUMBER: 120:226756

TITLE: Targeting delivery of protein drugs by chemical modification

AUTHOR(S): Hashida, Mitsuru; Nishikawa, Makiya; Yamashita, Fumiyoshi; Takakura, Yoshinobu

CORPORATE SOURCE: Fac. Pharm. Sci., Kyoto Univ., Kyoto, 606-01, Japan

SOURCE: Drug Dev. Ind. Pharm. (1994), 20(4), 581-90

CODEN: DDIPD8; ISSN: 0363-9045

DOCUMENT TYPE: Journal

LANGUAGE: English

AB In vivo disposition profiles of protein derivs. having various chem. modifications were systematically compared in mice based on the clearance concept. Proteins such as bovine .gamma.-globulin (IgG), bovine serum albumin (BSA), superoxide dismutase (SOD), soybean trypsin inhibitor (STI), and chicken egg white lysozyme (LZM) were (1) **conjugated** and **polyethylene glycol (PEG)** and dextran to increase mol. size, (2) **conjugated** with **carboxymethyl-dextran (CMD)** and diethylaminoethyl-dextran (DEAED) or coupled with diaminohexane or succinic acid to introduce elec. charges, and (3) modified with **galactose (Gal)** and mannose (Man) moieties to bestow an affinity for receptor-mediated endocytosis in cells. By applying these modifications, in vivo disposition features of proteins were extensively changed; i.e., in the case of SOD, **conjugation** with **CMD** and **PEG** prolonged its circulation half-life more than 100 times but cationized SOD showed remarkable accumulation on the surface of the liver tissue. In addn., specific targeting to the parenchymal cells of the liver was demonstrated in Gal-SOD, while, Man-SOD and succinylated SOD showed rapid uptake by the nonparenchymal cells. These results revealed the utility of chem. modification for controlling in vivo disposition of proteins.

IT 9001-63-2, Lysozyme 9054-89-1, Superoxide dismutase  
9078-38-0, Soybean trypsin inhibitor

RL: PROC (Process)

(chem. modification of, for targeting delivery)

RN 9001-63-2 HCAPLUS  
CN Lysózyme (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE


RN 9054-89-1 HCAPLUS  
CN Dismutase, superoxide (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9078-38-0 HCAPLUS  
CN Trypsin inhibitor, soybean (9CI) (CA INDEX NAME)


\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

IT **25322-68-3, Polyethylene glycol**  
RL: BIOL (Biological study)  
(protein drugs modified by, for targeting delivery)  
RN 25322-68-3 HCAPLUS  
CN Poly(oxy-1,2-ethanediyl), .alpha.-hydro-.omega.-hydroxy- (9CI) (CA INDEX NAME)



IT **59-23-4, Galactose, reactions 110-15-6, Succinic acid, reactions 124-09-4, Diaminohexane, reactions 3458-28-4, D-Mannose 9004-54-0, Dextran, reactions 9015-73-0, Diethylaminoethyl-dextran 9044-05-7, Carboxymethyl-dextran**  
RL: RCT (Reactant)  
(protein drugs modified by, for targeting delivery)  
RN 59-23-4 HCAPLUS  
CN D-Galactose (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



RN 110-15-6 HCAPLUS  
CN Butanedioic acid (9CI) (CA INDEX NAME)




RN 124-09-4 HCAPLUS  
CN 1,6-Hexanediamine (7CI, 8CI, 9CI) (CA INDEX NAME)



RN 3458-28-4 HCAPLUS  
CN D-Mannose (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).





RN 9004-54-0 HCAPLUS  
CN Dextran (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9015-73-0 HCAPLUS  
CN Dextran, 2-(diethylamino)ethyl ether (9CI) (CA INDEX NAME)


CM 1

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 100-37-8  
CMF C6 H15 N O



RN 9044-05-7 HCAPLUS  
CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)


CM 1

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
CMF C2 H4 O3



L47 ANSWER 18 OF 22 HCAPLUS COPYRIGHT 2002 ACS  
ACCESSION NUMBER: 1994:186083 HCAPLUS  
DOCUMENT NUMBER: 120:186083  
TITLE: Functional changes of lysozyme by **conjugating**  
with **carboxymethyl dextran**  
AUTHOR(S): Hattori, Makoto; Imamura, Shigeo; Nagasawa, Koichi;

CORPORATE SOURCE: Takahashi, Koji  
 Fac. Agric., Tokyo Univ. Agric. Technol., Tokyo, 183,  
 Japan  
 SOURCE: Biosci., Biotechnol., Biochem. (1994), 58(1), 174-7  
 CODEN: BBBIEJ; ISSN: 0916-8451  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Hen egg lysozyme-**carboxymethyl dextran** (HEL-  
**CMD**) **conjugate** was prepd. by using water-sol.  
 carbodiimide as a model protein-acidic **polysaccharide**  
**conjugate** for improving the protein function. An acid-amide bond  
 between HEL and **CMD** was confirmed by SDS-PAGE, isoelec. focusing  
 and IR spectra. The molar ratio of **CMD** to HEL in the  
**conjugate** was 1:1. The isoelec. point of the **conjugate**  
 was 5.5-6.0, which is much lower than that of HEL. Spectroscopic studies  
 suggested that the conformation around the Trp residue had not changed but  
 the .alpha.-helix content had decreased to about 1/3 that for native HEL.  
 The **conjugate** maintained about 60% of the enzymic activity of  
 native HEL at 40-60 .degree.C, while it was about 1.4 times as active as  
 native HEL at 4 .degree.C and 80 .degree.C. The **conjugate** was  
 more stable to proteolysis than native HEL. The denaturation temp. of the  
**conjugate** was about 73 .degree.C, which is almost the same as that  
 of native HEL. However, the enthalpy for denaturation of the  
**conjugate** was about 1/3 that of native HEL, which corresponds to  
 the decrease in .alpha.-helix content.

IT 9001-63-2D, Lysozyme, **conjugates** with  
**carboxymethyl dextran** 9044-05-7D,  
**Carboxymethyl dextran**, **conjugates** with  
 lysozyme

RL: BIOL (Biological study)  
 (conformation and thermal stability and catalytic properties of)

RN 9001-63-2 HCAPLUS

CN Lysozyme (8CI, 9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

RN 9044-05-7 HCAPLUS

CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)


CM 1

CRN 9004-54-0  
 CMF Unspecified  
 CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
 CMF C2 H4 O3



L47 ANSWER 19 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1993:678527 HCAPLUS

DOCUMENT NUMBER: 119:278527

searched by Susan Hanley 305-4053

TITLE: Synthesis and pharmacokinetics of a new liver-specific carrier, glycosylated carboxymethyl-dextran, and its application to drug targeting

AUTHOR(S): Nishikawa, Makiya; Kamijo, Akiko; Fujita, Takuya; Takakura, Yoshinobu; Sezaki, Hitoshi; Hashida, Mitsuru

CORPORATE SOURCE: Fac. Pharm. Sci., Kyoto Univ., Kyoto, 606-01, Japan

SOURCE: Pharm. Res. (1993), 10(9), 1253-61  
CODEN: PHREEB; ISSN: 0724-8741

DOCUMENT TYPE: Journal

LANGUAGE: English

AB To develop a new carrier system for hepatic targeting, **carboxymethyl-dextran (CMD)** was modified with **galactose** and mannose residues (**Gal-CMD**, **Man-CMD**), and their disposition characteristics were studied in mice using <sup>14</sup>C-labeled dextran. At a dose of 1 mg/kg, i.v.-injected **Gal-CMD** and **Man-CMD** rapidly accumulated in the liver parenchymal and nonparenchymal cells, resp., because of their preferential uptake via carboxylate receptors in these cells. Pharmacokinetic anal. revealed that their uptake rates were sufficiently large for selective drug targeting. Targeting of cytosine .beta.-D-arabinoside (araC) was studied using **Gal-CMD** as a sp. carrier to the hepatocytes. From the **conjugate** of araC with **Gal-CMD**, araC was released with a half-life of 36 h in phosphate buffer (pH 7.4) and 23 h in plasma. An in vivo biodistribution study demonstrated a disposition profile of the **conjugated** araC similar to that of the carrier, and selective delivery to hepatocytes of up to 80% of the dose was achieved. These findings suggest that glycosylated **CMDs** are carriers with a high affinity to liver parenchymal and nonparenchymal cells without any affinity to other tissues.

IT **9044-05-7**, Carboxymethyl dextran  
RL: RCT (Reactant)  
(glycosylation of, for drug targeting to liver)

RN 9044-05-7 HCAPLUS

CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 9004-54-0

CMF Unspecified


CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1

CMF C2 H4 O3



IT **151615-76-8P 151615-77-9P 151615-78-0P 151615-79-1P**  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(prepn. and pharmacokinetics of, for drug targeting to liver)


RN 151615-76-8 HCAPLUS

CN Dextran, 2-[(1-.beta.-D-arabinofuranosyl-1,2-dihydro-2-oxo-4-pyrimidinyl)amino]-2-oxoethyl carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 171340-32-2  
CMF C11 H15 N3 O7  
CDES 5:B-D-ARABINO

Absolute stereochemistry.




CM 2

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 3

CRN 79-14-1  
CMF C2 H4 O3




RN 151615-77-9 HCAPLUS  
CN Dextran, 2-[(1-.beta.-D-arabinofuranosyl-1,2-dihydro-2-oxo-4-pyrimidinyl)amino]-2-oxoethyl carboxymethyl 2-[[2-[[2-(.beta.-D-galactopyranosylthio)-1-iminoethyl]amino]ethyl]amino]-2-oxoethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 171340-34-4  
CMF C12 H23 N3 O7 S  
CDES 5:B-D-GALACTO


Absolute stereochemistry.



CM 2

CRN 171340-32-2  
CMF C11 H15 N3 O7  
CDES 5:B-D-ARABINO

Absolute stereochemistry.




CM 3

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 4

CRN 79-14-1  
CMF C2 H4 O3




RN 151615-78-0 HCAPLUS  
CN Dextran, carboxymethyl 2-[[2-[[1-imino-2-(.alpha.-D-mannopyranosylthio)ethyl]amino]ethyl]amino]-2-oxoethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 171340-33-3  
CMF C12 H23 N3 O7 S  
CDES 5:A-D-MANNO

Absolute stereochemistry.




CM 2

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 3

CRN 79-14-1  
CMF C2 H4 O3




RN 151615-79-1 HCAPLUS  
CN Dextran, carboxymethyl 2-[[2-[[2-(.beta.-D-galactopyranosylthio)-1-iminoethyl]amino]ethyl]amino]-2-oxoethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 171340-34-4  
CMF C12 H23 N3 O7 S  
CDES 5:B-D-GALACTO

Absolute stereochemistry.




CM 2

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN


\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 3

CRN 79-14-1  
CMF C2 H4 O3




IT **107-15-3**, 1,2-Ethanediamine, reactions  
 RL: RCT (Reactant)  
 (reaction of, with carboxymethyl dextran)  
 RN 107-15-3 HCAPLUS  
 CN 1,2-Ethanediamine (9CI) (CA INDEX NAME)




IT **75204-21-6 151530-12-0**  
 RL: RCT (Reactant)  
 (reaction of, with carboxymethyl dextran ethylenediamine deriv.)  
 RN 75204-21-6 HCAPLUS  
 CN Ethanimidic acid, 2-(.beta.-D-galactopyranosylthio)-, methyl ester (9CI)  
 (CA INDEX NAME)

Absolute stereochemistry.




RN 151530-12-0 HCAPLUS  
 CN D-Mannopyranose, 1-thio- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT **147-94-4**, Ara-C  
 RL: BIOL (Biological study)  
 (targeting of, to liver, glycosylated carboxymethyl dextran deriv. for)  
 RN 147-94-4 HCAPLUS  
 CN 2(1H)-Pyrimidinone, 4-amino-1-.beta.-D-arabinofuranosyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L47 ANSWER 20 OF 22 HCAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1993:116685 HCAPLUS

DOCUMENT NUMBER: 118:116685

TITLE: Therapeutic effects of superoxide dismutase derivatives modified with mono- or polysaccharides on hepatic injury induced by ischemia/reperfusion

AUTHOR(S): Fujita, Takuya; Furitsu, Hisao; Nishikawa, Mikiya; Takakura, Yoshinobu; Sezaki, Hitoshi; Hashida, Mitsuru

CORPORATE SOURCE: Fac. Pharm. Sci., Kyoto Univ., Kyoto, 606-01, Japan

SOURCE: Biochem. Biophys. Res. Commun. (1992), 189(1), 191-6

CODEN: BBRCA9; ISSN: 0006-291X

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Therapeutic effects of four types of recombinant superoxide dismutase (SOD) derivs., **conjugates** with polysaccharides, carboxymethyl (SOD-**CMD**) and diethylaminoethyl (SOD-DEAED) dextrans and galactosylated (Gal-SOD) and mannosylated (Man-SOD) derivs., on hepatic ischemia/reperfusion injury were studied in rats. Hepatic injury induced by transient occlusion and subsequent reflow of hepatic blood was evaluated by the anal. of biliary excretion of bromosulphophthalein (BSP) injected i.v. At a dose of 1000 units/kg, native SOD and SOD-DEAE had no significant effect and SOD-**CMD** had a slight effect. On the other hand, Gal-SOD and Man-SOD, targeted to liver parenchymal and nonparenchymal cells, resp., by a receptor-mediated endocytosis, exhibited superior inhibitory effects. These results demonstrated that these glycosylated SOD derivs. were useful for the prevention of hepatic ischemia/reperfusion injury.

IT **59-23-4D**, D-Galactose, superoxide dismutase **conjugates** **3458-28-4D**, D-Mannose, superoxide dismutase conjugates **9015-73-0D**, Diethylaminoethyl dextran, superoxide dismutase conjugates **9044-05-7D**, Carboxymethyldextran, superoxide dismutase conjugates **9054-89-1D**, Superoxide dismutase, polysaccharide derivs.


RL: BAC (Biological activity or effector, except adverse); BIOL (Biological study)

(antioxidant activity of, in liver injury from ischemia/reperfusion)

RN 59-23-4 HCAPLUS

CN D-Galactose (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




RN 3458-28-4 HCAPLUS

CN D-Mannose (9CI) (CA INDEX NAME)




Absolute stereochemistry. Rotation (+).



RN 9015-73-0 HCAPLUS  
 CN Dextran, 2-(diethylamino)ethyl ether (9CI) (CA INDEX NAME)  
 CM 1  
 CRN 9004-54-0  
 CMF Unspecified  
 CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE


CM 2  
 CRN 100-37-8  
 CMF C6 H15 N O



RN 9044-05-7 HCAPLUS  
 CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)  
 CM 1  
 CRN 9004-54-0  
 CMF Unspecified  
 CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2  
 CRN 79-14-1  
 CMF C2 H4 O3



RN 9054-89-1 HCAPLUS  
 CN Dismutase, superoxide (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

L47 ANSWER 21 OF 22 HCAPLUS COPYRIGHT 2002 ACS  
 ACCESSION NUMBER: 1993:66727 HCAPLUS  
 DOCUMENT NUMBER: 118:66727  
 TITLE: Targeted delivery of human recombinant superoxide  
 dismutase by chemical modification with mono- and

AUTHOR(S): polysaccharide derivatives  
Fujita, Takuya; Nishikawa, Makiya; Tamaki, Chieko;  
Takakura, Yoshinobu; Hashida, Mitsuru; Sezaki, Hitoshi  
CORPORATE SOURCE: Fac. Pharm. Sci., Kyoto Univ., Kyoto, 606-01, Japan  
SOURCE: J. Pharmacol. Exp. Ther. (1992), 263(3), 971-8  
CODEN: JPETAB; ISSN: 0022-3565

DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Four types of superoxide dismutase (SOD) derivs. such as SOD-**carboxymethyl dextran conjugate**, SOD-diethylaminoethyl dextran **conjugate**, galactosylated SOD and mannosylated SOD were synthesized and their potential for selective targeting to organs or cells was evaluated in mice by pharmacokinetic anal. All SOD derivs. retained 50 to 80% for 3 h. After i.v. injection, native SOD was rapidly excreted into urine and no significant accumulation was obsd. in the organs except the kidney. SOD-**carboxymethyl dextran conjugate** gave a long plasma half-life because of impaired glomerular filtration and tissue interaction. By contrast, galactosylated SOD and mannosylated SOD were very rapidly eliminated from the circulation and taken up by parenchymal and nonparenchymal cells of the liver, resp., via receptor-mediated endocytosis. These uptake processes were nonlinear and hepatic uptake clearance decreased as the dose increased, although almost complete extn. was obtained at a dose of 0.1 mg/kg. Furthermore, the accumulation in kidney of both glycosylated SODs was drastically decreased due to reduced renal proximal tubular reabsorption and also enhanced hepatic clearance. SOD-diethylaminoethyl dextran **conjugate** also rapidly disappeared from plasma and distributed into liver, but its accumulation occurred due to electrostatic interaction and was nonspecific in cellular distribution. These results suggest the possibility of controlling the in vivo fate of SOD at a cellular level by chem. modification utilizing sugar moieties with varied physicochem. and/or biol. characteristics.

IT **9054-89-1**, Superoxide dismutase  
RL: BIOL (Biological study)  
(human recombinant, targeted delivery of, by chem. modification with polysaccharides)

RN 9054-89-1 HCAPLUS

CN Dismutase, superoxide (9CI) (CA INDEX NAME)


\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

IT **59-23-4DP**, Galactose, **conjugates** with superoxide dismutase **3458-28-4DP**, Mannose, **conjugates** with superoxide dismutase **9015-73-0DP**, Diethylaminoethyl dextran, **conjugates** with superoxide dismutase **9044-05-7DP**, **Carboxymethyl dextran**, **conjugates** with superoxide dismutase **9054-89-1DP**, Superoxide dismutase, **conjugates** with polysaccharides  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(prepn. of, for targeted drug delivery)

RN 59-23-4 HCAPLUS

CN D-Galactose (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



RUSSEL 09/807,980

RN 3458-28-4 HCAPLUS  
CN D-Mannose (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



RN 9015-73-0 HCAPLUS  
CN Dextran, 2-(diethylamino)ethyl ether (9CI) (CA INDEX NAME)


CM 1

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 100-37-8  
CMF C6 H15 N O



RN 9044-05-7 HCAPLUS  
CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)


CM 1

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
CMF C2 H4 O3



RN 9054-89-1 HCAPLUS  
CN Dismutase, superoxide (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

L47 ANSWER 22 OF 22 HCAPLUS COPYRIGHT 2002 ACS  
ACCESSION NUMBER: 1985:571374 HCAPLUS

DOCUMENT NUMBER: 103:171374  
 TITLE: Carbohydrate-containing derivatives of the  
 trypsin-kallikrein inhibitor aprotinin from bovine  
 organs. II. Inhibitor coupled to the  
 (carboxymethyl)dextran derivatives of D-galactose  
 AUTHOR(S): Larionova, N. I.; Mityushina, G. V.; Kazanskaya, N.  
 F.; Blidchenko, Yu. A.; Berezin, I. V.  
 CORPORATE SOURCE: Dep. Chem., M. V. Lomonosov Moscow State Univ.,  
 Moscow, 119899, USSR  
 SOURCE: Biol. Chem. Hoppe-Seyler (1985), 366(8), 743-8  
 CODEN: BCHSEI  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB The trypsin-kallikrein inhibitor aprotinin was coupled to 2 ( **carboxymethyl)dextran** derivs. of D-galactose;  
 the **conjugates** contained 14 and 38 D-galactose  
 residues/mol of protein, resp. The apparent dissocn. consts.  $K_i$  of the  
 complexes between trypsin and modified aprotinins proved to be one order  
 of magnitude higher than the resp. values for native aprotinin. The  
 distribution of the modified aprotinins in rat organs after endocardial  
 injection was studied. The **conjugates** of aprotinin with ( **carboxymethyl)dextran** derivs. of D-galactose  
 were characterized by decreased clearance rates; they accumulated in the  
 active form in liver. The accumulation was 2.5-10 times higher than  
 native aprotinin for the time of observation (5 min-2 h).  
 IT **9044-05-7DP**, galactose derivs., aprotinin complexes  
**9087-70-1DP**, complexes with (carboxymethyl)dextran galactose  
 derivs.  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and pharmacokinetics of)  
 RN 9044-05-7 HCAPLUS  
 CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)


CM 1

CRN 9004-54-0  
 CMF Unspecified  
 CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1  
 CMF C2 H4 O3




RN 9087-70-1 HCAPLUS  
 CN Trypsin inhibitor, pancreatic basic (9CI) (CA INDEX NAME)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

IT **98913-51-0DP**, reaction products with lactose  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and reaction with lactose)  
 RN 98913-51-0 HCAPLUS  
 CN Dextran, 2-[(4-aminobutyl)amino]-2-oxoethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 171263-10-8  
CMF C6 H14 N2 O2



CM 2

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

IT **98913-51-0P**


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(prepn. and redn. of)

RN 98913-51-0 HCAPLUS

CN Dextran, 2-[(4-aminobutyl)amino]-2-oxoethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 171263-10-8  
CMF C6 H14 N2 O2



CM 2

CRN 9004-54-0  
CMF Unspecified  
CCI PMS, MAN


\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

IT **110-60-1**

RL: RCT (Reactant)  
(reaction of, with (carboxymethyl)dextran and carbodiimide)

RN 110-60-1 HCAPLUS

CN 1,4-Butanediamine (8CI, 9CI) (CA INDEX NAME)




IT **63-42-3**

RL: RCT (Reactant)  
(reaction of, with butylamino(carboxymethyl)dextran)

RN 63-42-3 HCAPLUS

CN D-Glucose, 4-O-.beta.-D-galactopyranosyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



IT 9044-05-7

RL: RCT (Reactant)

(reaction of, with diaminobutane and carbodiimide)

RN 9044-05-7 HCAPLUS

CN Dextran, carboxymethyl ether (9CI) (CA INDEX NAME)

CM 1

CRN 9004-54-0

CMF Unspecified

CCI PMS, MAN

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE

CM 2

CRN 79-14-1

CMF C2 H4 O3

